

Antenatal clinic HIV data found to underestimate actual prevalence declines: evidence from Zambia

Charles Michelo^{1,2}, Ingvild Sandøy² and Knut Fylkesnes²

1 Department of Community Medicine, School of Medicine, University of Zambia, Lusaka, Zambia

2 Centre for International Health, University of Bergen, Bergen, Norway

Summary

OBJECTIVE To determine to what extent antenatal clinic (ANC)-based estimates reflect HIV prevalence trends among men and women in a high prevalence urban population.

METHODS Examination of data from serial population-based HIV surveys in 1995 ($n = 2115$), 1999 ($n = 1962$) and 2003 ($n = 2692$), and ANC-based surveillance in 1994 ($n = 450$), 1998 ($n = 810$) and 2002 ($n = 786$) in the same site in Lusaka, Zambia. The population-based surveys recorded refusal rates between 6% and 10% during the three rounds.

RESULTS Among ANC attendees, prevalence declined by 20% (25.0% to 19.9%; $P = 0.101$) in the age group 15–24 years and was stable overall. In the general population, the prevalence declined by 49% ($P < 0.001$) and by 32% ($P < 0.001$) in age group 15–24 and 15–49, respectively. Among women only, HIV prevalence declined by 44% (22.5% to 12.5%; $P < 0.001$) and by 27% (29.6% to 21.7%; $P < 0.001$) in age group 15–24 and 15–49 years, respectively. In addition, prevalence substantially declined in higher educated women aged 15–24 years (20.7% to 8.5%, $P < 0.001$).

CONCLUSION ANC-based estimates substantially underestimated declines in HIV prevalence in the general population. This seemed to be partially explained by a combination of marked differentials in prevalence change by educational attainment and changes in fertility-related behaviours among young women. These results have important implications for the interpretation of ANC-based HIV estimates and underscore the importance of population-based surveys.

keywords HIV prevalence, antenatal, population survey, trends and Zambia

Introduction

HIV prevalence among pregnant women attending antenatal clinics (ANCs) remains the principal data source of infection trends in sub-Saharan Africa (Kwesigabo *et al.* 2000; Ghys *et al.* 2006). This type of surveillance has been revised to meet changing needs yet maintaining the original objective (WHO/UNAIDS 2003). Despite this usefulness, ANC-based HIV prevalence estimates should be interpreted with caution due to multiple potential inherent selection biases (Mills *et al.* 2005). Methods of adjustment for some of the factors have improved the estimation (Boisson *et al.* 1996; Zaba *et al.* 2000; Fabiani *et al.* 2003, 2006). Despite these drawbacks regarding point estimates, it has often been argued that ANC sentinel surveillance data collected over time will reasonably well capture infection trends of men and women in the general population (Kwesigabo *et al.* 2000). This argument is based on the assumption that the inherent biases remain constant over time, hence will not influence patterns

drastically (Mills *et al.* 2005). In the 1990s, very few communities had serial data to check this hypothesis and most validations of ANC-based HIV prevalence used single time-points or relatively short periods of time raising validity and accuracy concerns about the reliability of such extrapolations (Fylkesnes *et al.* 2001; Ghys *et al.* 2006).

In Zambia, the HIV epidemic has been monitored using both ANC-based and population-based data. The ANC-based surveillance system was established with few sites in 1990, and in 1994 a total of 27 sites were selected representing both rural and urban populations in all the provinces (Fylkesnes *et al.* 1998, 2001). Since then, three other rounds of surveillance have been conducted in 1998, 2002 and 2004 (Sandøy *et al.* 2006; Central Board of Health 2005). In one of the surveillance sites in Lusaka urban, serial cross-sectional surveys on HIV prevalence and risk factors were also conducted in 1995, 1999 and 2003 among randomly selected men and women (Fylkesnes *et al.* 2001; Michelo *et al.* 2006). Presently, this is the only site that has both ANC-based and population-based HIV

C. Michelo *et al.* Validating ANC-based HIV trends, Zambia

prevalence estimates from the same population consistently. We investigated how well antenatal-based HIV prevalence estimates capture trends in this general population between 1994 and 2003.

Methods**Antenatal-based surveillance design**

The data were collected from serial antenatal-based HIV surveys conducted in Chelstone in 1994 ($n = 450$), 1998 ($n = 810$) and 2002 ($n = 786$). The first epidemiological HIV sentinel surveillance among ANC attendees in Chelstone was conducted in 1990 as a pilot followed by another in 1993. However, the sample sizes in these surveys were very low. In 1994, the core antenatal-based HIV surveillance for the whole country was established, repeated in 1998 and in 2002. The detailed methods and major findings of the earlier national surveys have been reported elsewhere (Fylkesnes *et al.* 1998, 2001; Sandøy *et al.* 2006). Pregnant women who attended the ANC for the first time in the pregnancy were enrolled consecutively. Data were collected in a period not exceeding 4 months. Like in all the other major urban surveillance centres, the target number for Chelstone was 500 in 1994, but this was adjusted to 800 participants thereafter because of higher proportions of antenatal women in these sites compared with other and less urbanized areas. Serum from blood samples drawn for syphilis screening was tested unlinked and anonymously using Capillus HIV-1/HIV-2 rapid test (Cambridge Biotechnology, Galway, Ireland) at the ANCs. Randomly selected negative samples (5% in 1994 and 1998, 10% in 2002) and all positive samples were re-tested at the national laboratory using Wellcozyme HIV Recombinant HIV-1 (Murex, Johannesburg, South Africa). A third test, Bionor HIV-1 & 2 (Bionor AS, Skien, Norway), was employed on the samples with discordant results of tests one and two, and this third result was considered final.

Population-based surveillance design

The first population-based HIV survey in Zambia was conducted in 1995 in Chelstone and Kapiri Mposhi and two follow-up surveys were later conducted. The detailed methods and major findings of these population-based studies have been reported elsewhere (Fylkesnes *et al.* 1997, 1998, 2001; Michelo *et al.* 2006). The data used in this study were collected from the surveys conducted in Chelstone in 1995 ($n = 2115$), 1999 ($n = 1986$) and 2003 ($n = 2589$) using random cluster sample design described below. The Zambian census population mapping system was used to establish the sampling frame, which consisted

of 24 Standard Enumeration areas (SEAs) with 2786 households. Using 'probability proportional to size', 10 SEAs were selected for this study. In the sampled clusters, a personal structured interview was carried out with all eligible household members aged ≥ 15 years in order to collect information on education, socio-demographic characteristics and risk behaviours. The second part of the interview involved HIV testing using saliva. In 1995, all saliva samples were tested using Gacelisa HIV 1 & 2 (Welcome Diagnostics, Dartford, Kent, UK) and initially 450 randomly selected samples were tested using Bionor HIV-1 & 2 (Bionor AS) magnetic particle assay following modifications for saliva. Agreement between the two test kits was 99.8% (Fylkesnes & Kasumba 1998). Bionor 1 & 2 was used to test samples in the 1999 and 2003 follow-up surveys.

Validation strategy and statistical analyses

The ANC surveillance was conducted from August to November of 1994, 1998 and 2002. Each population survey was conducted within a year of the antenatal surveillance, that is, from October to December in 1995, October 1998 to May 1999 and from February to May in 2003. For comparability purposes, these years have been denoted as period 1, 2 and 3 to represent 1994, 1998 and 2002 for the ANC-based reports, whereas it is 1995, 1999 and 2003 in the population data, respectively. The health post where the ANC services are provided serves the same catchment area from which the sample for the population survey was drawn. Although in theory women in this area have a choice whether to go to another clinic for antenatal services or not, in practice this does not happen unless one is referred there (and vice versa) for a particular reason. ANC data from the 1990 pilot, 1993 and 2004 surveys were excluded from the analyses due to lack of appropriate population data for validation at the time. Analyses (stratified by age and sex) were performed using INTERCOOLED STATA version 8 (Texas, USA). The Mantel-Haenszel chi-square test (1 degree of freedom and with continuity correction) adjusted for cluster effect was used to test the linear trend of population-based HIV prevalence patterns over the periods. In order to check the effect of age structure, the estimates were standardized using the 2000 census urban female reference population. However, Clayton and Hills' score test for trend was employed for calculating linear trend in the ANC-based data. The median age of sexual debut was calculated using survival analysis and a log rank test for equality of survival curves was used to compare the median ages. To understand the possible differences and or similarities between ANC- and population-based respondents, further analyses were made in terms

C. Michelo *et al.* **Validating ANC-based HIV trends, Zambia**

of educational attainment, sexual debut and fertility patterns. The response to the question 'have you ever given birth?' was used to approximate fertility. The ANC-based estimates were then compared with the population-based estimates for similarity in all these aspects.

Ethics

The National AIDS Research Committee approved the protocol for the ANC-based surveillance system in 1990 and all HIV testing was done unlinked and anonymously as part of routine standard antenatal care in Zambia (Fylkesnes *et al.* 1998). The testing algorithm complied with the WHO/UNAIDS guidelines for conducting such surveys. The population-based survey protocols received clearance from the National AIDS Research Council and the University of Zambia Research Ethics Committee. The ethical details for both data sources are reported already elsewhere (Michelo *et al.* 2006; Sandøy *et al.* 2006).

Results

In the population-based survey, 60% of the respondents were females. The overall mean age was 26.2 years. The mean age among ANC attendees was 24.1 years. Furthermore, ANC attendees had significantly lower educational levels ($P < 0.001$) than respondents in the population-based survey (mean school years: 7.2 *vs.* 8.6 in period 1, 7.2 *vs.* 9.2 in period 2 and 7.6 *vs.* 10.1 in period 3). In the population-based surveys the overall response rate was 89%. Population-based antenatal attendance (during the last pregnancy) in the three surveys was 97%, 98% and 93% in 1995, 1999 and 2003, respectively.

Age-specific HIV infection trends

Table 1 illustrates both ANC and population-based age-specific HIV prevalence trends. Population-based HIV prevalence of men and women in age group 15–49 years between 1995 and 2003 declined by 32% (26.5%, 23.7% to 18.0%, $P < 0.001$). Among women aged 15–49, prevalence declined by 27% (29.6%, 27.1% to 21.7%, $P < 0.001$). In the same age group, ANC-based prevalence over the period remained stable (24.2%, 25.9% to 24.3%, $P = 0.821$). In age group 15–24 years, the population-based prevalence declined from 16.5%, 13.9% to 8.5% ($P < 0.001$) among men and women, representing a 49% decline and in young women only, prevalence declined by 44% (22.5%, 18.3% to 12.5%, $P < 0.001$). ANC-based data in this age group showed marginal decline of 20% (25.0%, 22.8% to 19.9%, $P = 0.101$). However, after adjusting for fertility risk, the ANC-based estimates

matched the estimates among young women aged 15–24 years in the general population. In this age group, fertility-adjusted prevalence declined by 42% (25.1%, 19.1% to 14.6%, $P = 0.001$) resembling the 44% decline observed in the general population of young women of the same age.

In the general population, prevalence declined largely in age group 15–29 years, whereas among ANC attendees, prevalence significantly declined only in the group aged 15–19 years. Furthermore, although ANC-based age-specific point estimates matched population-based estimates of young women aged 15–24 in time 1 (25% *vs.* 22.5%) and time 2 (22.8% *vs.* 18.3%), the ANC-based point estimate was significantly higher than population-based estimate in time 3 (19.9%, 95% CI 16.2–23.6.8 *vs.* 12.5%, 95% CI 9.2–15.8). When considering men and women together, ANC-based estimates still significantly over-estimated population prevalence in the groups aged 15–24 years throughout the survey rounds; time 1 (25%, 95% CI 19.8–30.2 *vs.* 16.5%, 95% CI 13.3–19.8), time 2 (22.8%, 95% CI 18.9–26.6 *vs.* 13.9%, 95% CI 12.3–15.5) and time 3 (19.9%, 95% CI 16.2–23.6 *vs.* 8.5%, 95% CI 6.5–10.6).

HIV and educational attainment

In general, population-based estimates showed a universal shift towards declining prevalence of HIV infection in groups with higher education over the period, as illustrated in Table 2. Among higher educated young men and women aged 15–24 years, prevalence declined from 16.1%, 11.7% to 5.3% ($P < 0.001$). In the same age group, prevalence declined from 20.7%, 16.1% to 8.5% ($P < 0.001$) among women only with higher education. Similarly, in higher educated groups of men and women aged 25–49 years, prevalence declined from 43.2% to 26.4% ($P < 0.001$) and from 45.6% to 29.0% ($P < 0.001$) in women only. Overall (men and women), prevalence declined from 32.0% to 14.8% ($P < 0.001$) and from 34.1% to 17.5% ($P < 0.001$) in women only. In sharp contrast, among ANC attendees, prevalence remained stable over the study in all age groups and irrespective of educational attainment. The overall level of educational attainment among ANC attendees did not significantly change over the periods (mean school years: 7.2, 7.2 and 7.6, $P = 0.219$; median school years 7.0 in all periods) despite an increase in the proportion of attendees with ≥ 11 school years from 10%, 13% to 21% in period 1, 2 and 3, respectively. Among young women aged 15–24 years, the pooled proportion of pregnant women with up to 7 years of school remained $>60\%$, whereas those with ≥ 11 school years was $<22\%$ in all the surveys. The distribution was the same even in age group 15–19 years.

C. Michelo *et al.* Validating ANC-based HIV trends, Zambia**Table 1** Age-specific adjusted prevalence differential trends of HIV infection by sex in Chelstone, Lusaka: 1995–2003

| Survey type | Time | Age-specific prevalence [% (95% CI*), n] | | | | | |
|---|------------------------------|--|---------------------------------|---------------------------------|---------------------------------|---------------------------------|----------------------------------|
| | | 15–19 | 20–24 | 25–29 | 30–39 | 40–49 | 15–49 |
| ANC sentinel surveillance | 1994 | 21.7% (13.7–29.7) n = 106 | 27.2% (20.2–34.1) n = 162 | 27.8% (18.9–37.2) n = 90 | 20.0% (11.3–28.7) n = 85 | | 24.2% (20.2–28.2) n = 450 |
| | 1998 | 15.1% (10.1–20.1) n = 199 | 28.4% (23.0–33.7) n = 275 | 31.9% (24.8–39.1) n = 166 | 29.4% (21.6–37.2) n = 136 | n = 7 | 25.9% (18.9–26.6) n = 474 |
| | 2002 | 9.2% (4.5–13.8) n = 153 | 25.5% (20.5–30.5) n = 294 | 28.7% (22.2–35.3) n = 188 | 33.1% (25.4–40.9) n = 145 | n = 10 | 24.3% (16.2–23.6) n = 447 |
| | Trend, P-value: 1994–2002 | 0.008 | 0.611 | 0.972 | 0.042 | – | 0.101 n = 786 |
| Population-based surveillance (females) | 1995 | 12.3% (7.4–17.1) n = 391 | 35.4% (26.7–43.9) n = 311 | 48.7% (42.6–54.9) n = 199 | 34.5% (28.0–41.0) n = 310 | 26.0% (13.8–38.2) n = 100 | 29.6% (26.8–32.4) n = 1311 |
| | 1999 | 9.5% (5.5–13.6) n = 336 | 27.9% (24.3–31.4) n = 305 | 41.2% (35.6–46.8) n = 221 | 39.3% (34.9–43.8) n = 244 | 22.3% (15.9–28.7) n = 112 | 27.1% (25.2–28.8) n = 1218 |
| | 2003 | 7.7% (4.7–10.6) n = 431 | 17.6% (13.3–21.9) n = 409 | 31.1% (26.1–36.1) n = 286 | 39.6% (33.9–45.2) n = 273 | 22.9% (17.0–28.9) n = 148 | 21.7% (19.4–24.1) n = 1547 |
| | Trend, P-value: 1995–2003 | 0.026 | <0.001 | <0.001 | 0.202 | 0.614 | <0.001 |
| Population-based surveillance (males and females) | 1995 | 9.1% (5.3–12.9) n = 657 | 26.7% (19.6–33.8) n = 479 | 43.1% (37.9–48.4) n = 313 | 36.3% (31.1–41.5) n = 460 | 33.9% (25.8–42.2) n = 206 | 26.5% (23.9–28.9) n = 2115 |
| | 1999 | 8.2% (5.2–11.2) n = 560 | 20.1% (17.6–23.6) n = 513 | 34.8% (29.7–40.2) n = 330 | 39.3% (34.9–43.6) n = 365 | 29.9% (24.7–35.1) n = 194 | 23.7% (22.3–25.2) n = 1961 |
| | 2003 | 5.4% (3.4–7.3) n = 745 | 11.8% (9.0–14.6) n = 718 | 24.7% (21.1–28.2) n = 446 | 37.9% (31.9–43.8) n = 454 | 26.5% (17.2–35.9) n = 226 | 18.0% (15.3–20.8) n = 2589 |
| | Trend, P-value: 1995–2003 | 0.007 | <0.001 | <0.001 | 0.626 | 0.091 | <0.001 |

Notes: 1, *Confidence Intervals adjusted for clustering effect in the population-based estimates; 2, chi-square for linear trend (all significant P-values are highlighted in bold); 3, sample sizes were (a) ANC: 450, 786 and 786 in 1994, 1998 and 2002, respectively (b) Population-based: 2115, 1961 and 2589 in 1995, 1999 and 2003, respectively.

C. Michelo *et al.* Validating ANC-based HIV trends, Zambia**Table 2** Comparing trends of HIV prevalence by years of schooling between antenatal clinic (ANC) attendees and the general population in (1995–2003) Chelstone

| Age | Population | Education (years) | Prevalence, % (<i>n</i>) | | | § MH Chi-square for linear trend |
|-------|---------------|-------------------|----------------------------|-----------------|------------------|----------------------------------|
| | | | Time 1 | Time 2 | Time 3 | Time 1 to 3 |
| 15–24 | ANC attendees | 0–7 | 27.2% (53/195) | 22.6% (56/248) | 20.8% (49/236) | 0.118 |
| | | 8–10 | 19.4% (7/36) | 21.0% (26/124) | 19.2% (23/120) | 0.840 |
| | | ≥11 | 20.0% (4/20) | 23.7% (14/59) | 18.5% (15/81) | 0.634 |
| | PBS (F) | 0–7 | 21.5% (59/274) | 18.4% (32/174) | 20.2% (35/173) | 0.664 |
| | | 8–10 | 23.8% (66/277) | 20.0% (50/250) | 13.8% (33/240) | 0.000 |
| | | ≥11 | 20.7% (30/145) | 16.1% (35/217) | 8.5% (36/423) | 0.000* |
| | PBS (M + F) | 0–7 | 16.8% (64/380) | 17.5% (41/234) | 15.9% (39/244) | 0.495 |
| | | 8–10 | 16.2% (77/476) | 14.0% (59/421) | 10.0% (44/440) | 0.011 |
| | | ≥11 | 16.1% (44/273) | 11.7% (49/418) | 5.3% (41/774) | 0.001* |
| 25–49 | ANC attendees | 0–7 | 26.2% (28/107) | 34.9% (51/146) | 29.8% (45/151) | 0.646 |
| | | 8–10 | 14.3% (3/21) | 27.4% (20/73) | 32.2% (19/59) | 0.166 |
| | | ≥11 | 21.1% (4/19) | 27.3% (9/33) | 33.8% (25/74) | 0.264 |
| | PBS (F) | 0–7 | 28.5% (63/221) | 26.7% (47/176) | 34.6% (55/159) | 0.288 |
| | | 8–10 | 41.2% (86/209) | 42.4% (86/203) | 36.5% (78/214) | 0.389 |
| | | ≥11 | 45.6% (77/169) | 39.9% (79/198) | 29.0% (96/331) | 0.001* |
| | PBS (M + F) | 0–7 | 30.1% (84/279) | 26.5% (56/211) | 36.1% (74/205) | 0.278 |
| | | 8–10 | 38.8% (116/299) | 38.2% (100/262) | 34.9% (98/281) | 0.213 |
| | | ≥11 | 43.2% (168/389) | 38.6% (160/415) | 26.4% (168/637) | 0.000* |
| 15–49 | ANC attendees | 0–7 | 26.8% (84/314) | 27.2% (116/426) | 24.3% (101/415) | 0.423 |
| | | 8–10 | 17.2% (10/58) | 22.5% (46/204) | 22.9% (44/192) | 0.469 |
| | | ≥11 | 22.0% (9/41) | 26.6% (25/94) | 26.1% (43/165) | 0.707 |
| | PBS (F) | 0–7 | 24.7% (122/495) | 22.6% (79/350) | 27.1% (90/332) | 0.657 |
| | | 8–10 | 31.3% (152/486) | 30.0% (136/453) | 24.5% (111/454) | 0.009 |
| | | ≥11 | 34.1% (107/314) | 27.5% (114/415) | 17.5% (132/754) | 0.000* |
| | PBS (M + F) | 0–7 | 22.5% (148/659) | 21.8% (97/445) | 25.2% (113/449) | 0.501 |
| | | 8–10 | 24.9% (193/775) | 23.3% (159/683) | 19.7% (142/721) | 0.033 |
| | | ≥11 | 32.0% (212/662) | 25.1% (209/833) | 14.8% (209/1411) | 0.000* |

Notes: 1. Years: ANC, 1 = 1994, 2 = 1998, 3 = 2002; PBS: 1 = 1995, 2 = 1999, 3 = 2003. 2. Sample sizes: PBS: *n* = 2115 in 1995; *n* = 1961 in 1999; *n* = 2589 in 2003; ANC: *n* = 450 in 1994; *n* = 786 in 1998; *n* = 786 in 2002. 3. Chi-square test for trend (a) § MH denotes Mantel–Haenszel chi square for linear trend adjusted for cluster effect (1 degree of freedom and with continuity correction), in the population-based estimates. However, Clayton and Hills' score test for trend was used in the ANC data (b) statistically significant *P*-values are highlighted in bold (c) * signifies *P* < 0.001.

Population-based sexual debut and ever given birth by educational attainment

In the general population of women aged 15–24 years, the mean number of years in school increased from 8.64 (95% CI 8.64–8.83) in 1995 and 9.21 (95% CI 9.04–9.38) in 1999 to 10.1 (95% CI 9.95–10.26) in 2003. In this group, older age at first sex was associated with higher educational attainment. In 2003, the median age for sexual debut among women aged 20–29 years and with >11 years of school was 19 years whereas it was 16.6 years in groups with 0–7 years of school (*P* < 0.001). Similarly, the proportion of young people aged 15–24 who said they ever had sex dropped between 1999 and 2003 (males: Adjusted

Odds Ratio (AOR) 0.83 95% CI 0.62–1.10; females: AOR 0.68 95% CI 0.52–0.88). Following a similar pattern, higher educated young people showed significant postponement in ages at first birth. Consequently, among women with >11 school years, the proportion ever given birth in age group 15–24 years decreased from 33.1%, 22.2% to 19.2% (*P* = 0.002) whereas the decline was marginal in groups with 0–7 school years (Figure 1). In age group 15–19, the proportion of women ever given birth declined by 57% (16.8%, 9.4% to 7.9%; OR 0.43, 95% CI 0.26–0.68) and from 65.5%, 47.5% to 41.7% (*P* < 0.000) in age group 20–24 years (see Table 3). Figure 2 illustrates further the general decline in age at first birth among women <30 years of age in the general population.


C. Michelo *et al.* Validating ANC-based HIV trends, Zambia

Figure 1 Proportion (per cent) of women 'ever given birth' by educational level in aged 15–24 years in Chelstone (urban), Zambia. Notes: 1, Sample sizes $n = 886, 853$ and 1018 in 1995, 1999 and 2003, respectively; 2, P -value for MH chi square for trend: 0–7 school years, $P = 0.405$, 8–10 school years, $P = 0.289$ and for ≥ 11 school years, $P = 0.0001$.

Discussion

We found diminishing representativeness of ANC-based data in capturing HIV prevalence and trends in the general population. First, the overall ANC-based HIV prevalence estimates remained stable, contrasting the population-based pattern where prevalence declined in both women and men. Second, both sources of data showed HIV declines in age group 15–24 years, but declines were substantially steeper in population-based data (which showed a decline of 49%) compared with ANC-based data (which showed a decline of 20%). These findings seem to be partially explained by a combination of marked differentials in prevalence change by educational attainment as well as marked changes in fertility-related behaviours such as the postponement of first birth among young women. This postponement was associated with educational attainment. We have already reported substantial HIV declines among higher educated groups in this population (Michelo *et al.* 2006). The




Figure 2 Increase in age at first birth (1995–2003) among HIV negative women ever given birth in Chelstone (Lusaka urban), Zambia. Notes: 1, Sample sizes in group aged 15–30 years, $n = 1511, 1462$ and 1994 in 1995, 1999 and 2003, respectively (overall all ages, $n = 6665$); 2, P -value for the increase in age at first birth was significant ($P < 0.05$) in all age groups < 31 years of age.

findings are a challenge to the interpretation of results from HIV surveillance systems, and whether the results are generalizable to other areas in Zambia and countries in sub-Saharan Africa merits further study. A similar pattern of change in ANC-based HIV prevalence reported here has been observed in the majority of surveillance sites in Zambia, and the most likely scenario is that similar change mechanisms were seen in many of these populations (Sandøy *et al.* 2006). Historically, the only and dramatic parallel declines and not associated with postponement of birth, observed from both antenatal- and population-based data was in Kagera region during the 1990s (Asiimwe-Okiror *et al.* 1997; Kwesigabo *et al.* 1998, 2000, 2005). The declines in Kagera may have been a signal of continuing declines in the general population as the epidemic matured. At such a stage of the epidemic as currently observed, dynamics among antenatal attendees differ from the general population; hence, using ANC-based estimates alone could even lead to a possibility of serious

Table 3 Population-based declining trend in the proportion of women ever given birth in Chelstone (urban), Zambia: 1995–2003

| Age | Proportion of women ever given birth in respective periods | | | P -value (trend‡) |
|-------|--|-----------------|-----------------------------------|---------------------|
| | 1995*, % (n) | 1999, % (n) | 2003, % (n); OR† 95% CI | |
| 15–19 | 16.8 (382) | 9.4 (330) | 7.9 (355); 0.43 95% CI 0.26–0.68 | <0.001 |
| 20–24 | 65.5 (310) | 47.5 (299) | 41.7 (384); 0.38 95% CI 0.28–0.51 | <0.001 |
| 25–29 | 85.4 (198) | 83.9 (218) | 74.8 (282); 0.51 95% CI 0.32–0.82 | 0.003 |
| 30–39 | 95.2 (310) | 95.0 (242) | 94.5 (273); 0.38 95% CI 0.28–0.51 | 0.722 |

*Sample sizes: $n = 1100, 1089$ and 1294 in 1995, 1999 and 2003 respectively.

†OR denotes odds ratio for the proportions in 2003 using the proportions in 1995 as a reference category and the 95% CI have adjusted for cluster effect.

‡Denotes Mantel–Haenszel chi square for linear trend adjusted for cluster effect (1 degree of freedom and with continuity correction).

C. Michelo *et al.* **Validating ANC-based HIV trends, Zambia**

underestimation of what could be intervention-linked impacts.

The basis for our assessment of how well ANC-based data capture HIV trends in the general population was data from a population-based survey conducted in the area covered by the ANC. There is a potential of non-response to have biased the results. Among ANC attendees, non-response can be estimated largely by ANC coverage, and coverage had been stable and above 90% in all the surveys (estimated to be 98% in Lusaka) (Central Statistical Office (Zambia) *et al.* 2003). In addition, movements of nearby populations of women to and from the ANC catchment area were the other possible sources of bias. However, the influence exerted by nearby populations on ANC prevalence estimates in our study was minimal as movement between ANCs to and from this catchment area is not by choice but dictated by a need-driven functioning referral system. One such need is the availability of Prevention of Mother to Child Transmission (PMTCT) services. In Chelstone, PMTCT services were only firmly grounded in 2004 and are unlikely to have been a factor in the stability of the ANC-based prevalence over the period in question. In the population-based study, the most significant cause of non-participation was absence of men. Refusal to give saliva for HIV testing was low in all the three surveys. In view of this it does not seem likely that non-response bias was an important factor affecting the results. In addition, fertility-linked factors which could be the other possible sources of selection bias were critical explanatory factors in this study. Furthermore, we also exclude laboratory-associated differentials in test result to have been a factor (Fylkesnes & Kasumba 1998; Fylkesnes *et al.* 1998).

There is convincing evidence that HIV transmission among adults in sub-Saharan Africa is predominantly heterosexual (Schmid *et al.* 2004). Women tend to under-report socially undesirable sexual behaviours, such as young age at sexual debut (Upchurch *et al.* 2002; Central Statistical Office (Zambia) *et al.* 2003). However, elsewhere we have argued that 'it is reasonable to believe that asking a woman whether she has ever given birth will give more reliable answers than whether she has ever had sex, as childbearing is associated with high respect in this society and is difficult to keep secret' (Sandøy *et al.* 2007). Therefore, the reduction in the proportion who had ever given birth and parallel postponement of age at first birth among higher educated young people in the general population suggests a convincing behaviour change. Postponement of childbearing might be due to a combination of abstinence, consistent use of condoms as well as utilization of other contraceptives, all of which have increased during the period (Central Statistical Office (Zambia) *et al.* 2003). These factors were of critical

importance in explaining the reduced odds of HIV infection among higher educated groups in the general population of young females (Fylkesnes *et al.* 1998; Michelo *et al.* 2006).

However, the picture is dissimilar among ANC attendees. First, the overestimation of HIV prevalence seen in ANC data from age group 15–19 years might be related to the fact that this group engaged in unprotected sex, thereby increasing the likelihood of infection compared with their counterparts in the general population, some of whom are not sexually active (Konde-Lule *et al.* 1997; Fylkesnes *et al.* 1998; Gray *et al.* 1998; Pettifor *et al.* 2004). Second, the higher prevalence in age group 20–24 years also suggests that there is a steady increase in the number of newly infected women arising from younger age groups. Since male HIV prevalence in parallel ages in the general population is generally low, there was evidence suggesting that these young women most probably were involved in cross-generation relationships and had unprotected sex with infected older men (Gregson *et al.* 2002). This could be the major reason why young antenatal women have a higher likelihood of HIV infection (Strickler *et al.* 1995; Fylkesnes *et al.* 1998; Fabiani *et al.* 2003; Mills *et al.* 2005). We also noted that the observed parallel HIV declines associated with education in the general population was absent among ANC attendees. Higher educated young women in the general population were postponing pregnancy and the postponement was substantial in a relatively short period of time. This is likely to be due to a combination of an ongoing process of fertility change which is probably linked with the success of other preventive strategies against HIV transmission such as condom use, abstinence and a longer and more careful search of 'safer' marital partners. In contrast, young ANC women may have ignored messages about abstinence and protection against HIV infection, which was probably influenced by biological and social pressures including the mere desire for motherhood. This differentiating feature between ANC attendees and young women in the general population merits further study and monitoring, especially since the population distribution continues to be highly dynamic.

Therefore, the interpretation of ANC-based HIV prevalence estimates and their extrapolation to the general population remain difficult tasks (Strickler *et al.* 1995; Mills *et al.* 2005). This is largely because in general, ANC-generated prevalence is also vulnerable to several time-dependent sources of bias such as usage and coverage of ANC services, migration, deaths, convenience sampling of sites, population distribution by age, differential distribution by social classes, contraceptive use and

C. Michelo *et al.* **Validating ANC-based HIV trends, Zambia**

fertility differences with HIV negative women including behavioural differences. This limits generalizability (Mills *et al.* 2005). Adjusting for fertility risk using data from a reference general population within the region improves estimates from ANC attendees (Fabiani *et al.* 2003). In our study, we achieved similar findings by adjusting for fertility risk among ANC attendees using the same source general population as a reference population.

In summary, our results have some important implications for the interpretation of ANC-based HIV estimates. They underscore the importance of population-based surveys, and particularly surveys conducted in selected communities in order to improve the interpretation of HIV trends captured by ANC attendees. These surveys should measure biological, behavioural and socio-demographic information concomitantly to generate critical knowledge for improving our understanding of dynamics of population responses. This also calls for adjustments in the current ANC surveillance systems to allow for sampling adjustments, collection of additional important details such as testing and treatment history as well as indicators for behavioural patterns, in order as to assess the extent of inherent biases which are in turn useful in interpreting ANC-based HIV trends.

Acknowledgements

The authors would like to thank the Chelstone participants, research assistants and staff at the Kabwe General Hospital, University Teaching Hospital virology laboratory unit, Central Statistical Office as well as the Zambia National AIDS/STD/TB & Leprosy programme (forerunner to the Zambia National HIV/AIDS/STI/TB Council). Funding was provided by the Norwegian government through the Norwegian Council for Higher Education's programme for Development Research and Education (NUFU) and the Norwegian Agency for Development Co-operation (NORAD).

References

- Asiimwe-Okiror G, Opio AA, Musinguzi J *et al.* (1997) Change in sexual behaviour and decline in HIV infection among young pregnant women in urban Uganda. *AIDS* 11, 1757–1763.
- Boisson E, Nicoll A, Zaba B & Rodrigues LC (1996) Interpreting HIV seroprevalence data from pregnant women. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology* 13, 434–439.
- Central Statistical Office, Central Board of Health (Zambia) & ORC Macro (2003) *Zambia Demographic and Health Survey 2001–2002*. ORC Macro, Calverton, MD.
- Fabiani M, Fylkesnes K, Nattabi B, Ayella EO & Declich S (2003) Evaluating two adjustment methods to extrapolate HIV prevalence from pregnant women to the general female population in sub-Saharan Africa. *AIDS* 17, 399–405.
- Fabiani M, Nattabi B, Ayella EO, Ogwang M & Declich S (2006) Differences in fertility by HIV serostatus and adjusted HIV prevalence data from an antenatal clinic in northern Uganda. *Tropical Medicine & International Health* 11, 182–187.
- Fylkesnes K & Kasumba K (1998) The first Zambian population-based HIV survey: saliva-based testing is accurate and acceptable. *AIDS* 12, 540–541.
- Fylkesnes K, Musonda RM, Kasumba K *et al.* (1997) The HIV epidemic in Zambia: socio-demographic prevalence patterns and indications of trends among childbearing women. *AIDS* 11, 339–345.
- Fylkesnes K, Ndhlovu Z, Kasumba K, Mubanga Musonda R & Sichone M (1998) Studying dynamics of the HIV epidemic: population-based data compared with sentinel surveillance in Zambia. *AIDS* 12, 1227–1234.
- Fylkesnes K, Musonda RM, Sichone M *et al.* (2001) Declining HIV prevalence and risk behaviours in Zambia: evidence from surveillance and population-based surveys. *AIDS* 15, 907–916.
- Ghys PD, Kufa E & George MV (2006) Measuring trends in prevalence and incidence of HIV infection in countries with generalised epidemics. *Sexually Transmitted Infections* 82(Suppl. 1), i52–i56.
- Gray RH, Wawer MJ, Serwadda D *et al.* (1998) Population-based study of fertility in women with HIV-1 infection in Uganda. *Lancet* 351, 98–103.
- Gregson S, Nyamukapa CA, Garnett GP *et al.* (2002) Sexual mixing patterns and sex-differentials in teenage exposure to HIV infection in rural Zimbabwe. *Lancet* 359, 1896–1903.
- Konde-Lule JK, Sewankambo N & Morris M (1997) Adolescent sexual networking and HIV transmission in rural Uganda. *Health Transition Review* 7(Suppl.), 89–100.
- Kwesigabo G, Killewo J, Godoy C *et al.* (1998) Decline in the prevalence of HIV-1 infection in young women in the Kagera region of Tanzania. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology* 17, 262–268.
- Kwesigabo G, Killewo JZ, Urassa W *et al.* (2000) Monitoring of HIV-1 infection prevalence and trends in the general population using pregnant women as a sentinel population: 9 years experience from the Kagera region of Tanzania. *Journal of Acquired Immune Deficiency Syndrome* 23, 410–417.
- Kwesigabo G, Killewo J, Urassa W *et al.* (2005) HIV-1 infection prevalence and incidence trends in areas of contrasting levels of infection in the Kagera region, Tanzania 1987–2000. *Journal of Acquired Immune Deficiency Syndrome* 40, 585–591.
- Michelo C, Sandøy IF & Fylkesnes K (2006) Marked HIV prevalence declines in higher educated young people: evidence from population-based surveys (1995–2003) in Zambia. *AIDS* 20, 1031–1038.

C. Michelo *et al.* **Validating ANC-based HIV trends, Zambia**

- Mills S, Rehle T & Schwartländer B (2005) Understanding HIV epidemics and the response to them: linking data on behaviour change and HIV decline. In: *Evaluating Programs for HIV/AIDS Prevention and Care in Developing Countries – A Handbook for Program Managers and Decision Makers, Vol. 2005* (eds T Rehle, T Saidel, S Mills & R Magnani). Family Health International, Arlington, VA, pp. 211–212.
- Pettifor AE, van der Straten A, Dunbar MS, Shiboski SC & Padian NS (2004) Early age of first sex: a risk factor for HIV infection among women in Zimbabwe. *AIDS* **18**, 1435–1442.
- Sandøy IF, Michelo C, Kvåle G & Fylkesnes K (2006) Antenatal clinic-based HIV prevalence in Zambia: declining trends but sharp local contrasts in young women. *Tropical Medicine & International Health*, **11**, 917.
- Sandøy IF, Michelo C, Siziya S & Fylkesnes K (2007) Associations between sexual behaviour change in young people and decline in HIV prevalence in Zambia. *BMC Public Health* **7**, 60.
- Schmid GP, Buve A, Mugenyi P *et al.* (2004) Transmission of HIV-1 infection in sub-Saharan Africa and effect of elimination of unsafe injections. *Lancet* **363**, 482–488.
- Strickler H, Hoover DR & Dersimonian R (1995) Problems in interpreting HIV sentinel seroprevalence studies. *Annals of Epidemiology* **5**, 447–454.
- Upchurch DM, Lillard LA, Aneshensel CS & Fang Li N (2002) Inconsistencies in reporting the occurrence and timing of first intercourse among adolescents. *Journal of Sex Research* **39**, 197–206.
- WHO/UNAIDS (2003) *Guidelines for Conducting HIV Sentinel Serosurveys Among Pregnant Women and Other Groups*. Joint United Nations Programme on HIV/AIDS, Geneva.
- Zaba BW, Carpenter LM, Boerma JT *et al.* (2000) Adjusting ante-natal clinic data for improved estimates of HIV prevalence among women in sub-Saharan Africa. *AIDS* **14**, 2741–2750.

Corresponding Author Charles Michelo, Department of Community Medicine, University of Zambia, PO Box 50110, Lusaka, Zambia. Tel.: +260 966 754920; Fax: +260 1 256181; E-mail: ccmichelo@yahoo.com