

# Progress in Physical Geography

<http://ppg.sagepub.com/>

---

## **Recent progress in understanding marine-terminating Arctic outlet glacier response to climatic and oceanic forcing: Twenty years of rapid change**

J. Rachel Carr, Chris R. Stokes and Andreas Vieli

*Progress in Physical Geography* published online 5 April 2013

DOI: 10.1177/0309133313483163

The online version of this article can be found at:

<http://ppg.sagepub.com/content/early/2013/04/05/0309133313483163>

---

Published by:



<http://www.sagepublications.com>

**Additional services and information for *Progress in Physical Geography* can be found at:**

**Email Alerts:** <http://ppg.sagepub.com/cgi/alerts>

**Subscriptions:** <http://ppg.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

>> [OnlineFirst Version of Record](#) - Apr 5, 2013

[What is This?](#)



# Recent progress in understanding marine-terminating Arctic outlet glacier response to climatic and oceanic forcing: Twenty years of rapid change

**J. Rachel Carr**

Durham University, UK

**Chris R. Stokes**

Durham University, UK

**Andreas Vieli**

Durham University, UK

## Abstract

Until relatively recently, it was assumed that Arctic ice masses would respond to climatic/oceanic forcing over millennia, but observations made during the past two decades have radically altered this viewpoint and have demonstrated that marine-terminating outlet glaciers can undergo dramatic dynamic change at annual timescales. This paper reviews the substantial progress made in our understanding of the links between marine-terminating Arctic outlet glacier behaviour and the ocean-climate system during the past 20 years, when many ice masses have rapidly lost mass. Specifically, we assess three primary climatic/oceanic controls on outlet glacier dynamics, namely air temperature, ocean temperature and sea ice concentrations, and discuss key linkages between them. Despite recent progress, significant uncertainty remains over the response of marine-terminating outlet glaciers to these forcings, most notably: (1) the spatial variation in the relative importance of each factor; (2), the contribution of glacier-specific factors to glacier dynamics; and (3) the limitations in our ability to accurately model marine-terminating outlet glacier behaviour. Our present understanding precludes us from identifying patterns of outlet glacier response to forcing that are applicable across the Arctic and we underscore the potential danger of extrapolating rates of mass loss from a small sample of study glaciers.

## Keywords

Arctic, climate change, cryosphere, Greenland Ice Sheet, marine-terminating outlet glaciers

## 1 Introduction


Arctic warming is expected to far exceed the global average and is forecast to reach 4–7°C by 2100 (IPCC, 2007; Meier et al., 2007). Consequently, Arctic ice masses are expected to

---

### Corresponding author:

J. Rachel Carr, Durham University, Science Laboratories, South Road, Durham DH1 3LE, UK.

Email: j.r.carr@durham.ac.uk




**Figure 1.** Regional overview map showing the location of major ice masses and other sites discussed in the text. Major water masses are also labelled. Glacier abbreviations are as follows: Helheim Glacier (HH), Kangerdlugssuaq Glacier (KG), Dugaard Jensen Gletscher (DJ), Kangerdlugssuaq Nunata Sermia (KNS), Jakobshavn Isbrae (JI), Petermann Glacier (PG), Hansbreen (HB), Duvebreen (DB) and John Evans Glacier (JEG). Inset: Overview map of Alaska, showing the location of LeConte Glacier (LCG).

undergo rapid change during the 21st century and to contribute significantly to global sea level rise (e.g. Bamber et al., 2007). Indeed, estimates suggest that the Greenland Ice Sheet (GIS) contributed  $0.46 \text{ mm a}^{-1}$  to sea level rise between 2000 and 2008 (van den Broeke et al., 2009). Assessing the potential response of Arctic ice masses to climate change is therefore crucial for the accurate prediction of near-future sea level rise (IPCC, 2007). For the purposes of this paper, we define ‘Arctic ice masses’ as the major glaciated archipelagos within the Arctic Circle, namely the Greenland Ice Sheet (GIS), Svalbard, Novaya Zemlya (NZ), Severnaya Zemlya (SZ), Franz Josef Land (FJL) and the Canadian Arctic (Figure 1).

Alaska is also included as results from the region have contributed significantly to our knowledge of marine-terminating outlet glacier dynamics. Here we define a marine-terminating outlet glacier as a channel of fast-moving ice that drains an ice cap or ice sheet and terminates in the ocean, at either a floating or grounded margin (Benn and Evans, 2010) (Figure 2).

Our understanding of Arctic ice mass behaviour has advanced dramatically during the last 20 years, particularly during the last decade. Previously, it was generally assumed that large Arctic ice masses would respond to climatic warming at millennial timescales, primarily through increased surface melting, and that changes in ice flow would occur only at



**Figure 2.** Visible satellite imagery of selected marine-terminating Arctic outlet glaciers and Arctic ice masses at 1:1,000,000 scale. Images are ordered by glacier location, from west to east, and show (A) Petermann Glacier, northwest Greenland; (B) Kangerdlugssuaq Glacier, east Greenland; (C) Vestfonna Ice Cap, Svalbard; (D) northern ice cap, Novaya Zemlya. Outlet glacier and ice mass locations are shown in Figure 1. Major outlet glaciers are labelled according to terminus type (M = marine; L = land) and approximate near-terminus flow direction is marked (dashed lines).

*Imagery source:* Global Land Cover Facility ([www.landcover.org](http://www.landcover.org)).

centennial timescales or longer (Greve, 2000; Huybrechts et al., 1991; IPCC, 2001). However, studies published during the past two decades have dramatically altered this viewpoint (e.g.

Joughin et al., 2010; Rignot et al., 2008; van den Broeke et al., 2009) and have shown that most Arctic ice masses have rapidly lost mass since the 1990s. Crucially, losses have been concentrated




at the coastal margins, particularly on marine-terminating outlet glaciers (e.g. Joughin et al., 2010; Meier et al., 2007; Thomas et al., 2009). Indeed, recent studies have demonstrated that marine-terminating Arctic outlet glaciers can respond rapidly to climatic/oceanic forcing (e.g. Andersen et al., 2012; Howat et al., 2007, 2008a, 2011; Joughin et al., 2008b, 2010; Kjær et al., 2012) and can significantly influence the mass budget of their parent ice masses over annual to decadal timescales (e.g. Pritchard et al., 2009; Rignot et al., 2008; Stearns and Hamilton, 2007).

Results from the Antarctic, particularly Pine Island Glacier (Payne et al., 2004), have also highlighted the role of outlet glaciers and ice streams in enabling rapid coupling between forcing at the margins and the ice sheet interior and have raised concerns over the vulnerability of some regions to rapid mass loss (Joughin and Alley, 2011). Furthermore, iceberg-rafted debris from palaeo-ice sheets attests to major episodes of ice sheet collapse (e.g. Bond et al., 1992) and reconstructions of marine-based palaeo-ice sheets have highlighted the potential for rapid ice stream/outlet glacier retreat (e.g. Briner et al., 2009; Winsborrow et al., 2010). Theoretical considerations also suggest that glaciers resting on reverse bed slopes may potentially be unstable (Thomas, 1979; Weertman, 1974). Although this review focuses on the Arctic, these findings have demonstrated that marine-terminating outlet glaciers can respond rapidly to climatic/oceanic forcing and play a key role in regulating the mass balance of marine-based ice sheets. As a result, the factors controlling marine-terminating outlet glacier dynamics have emerged as a primary area of research.

Recent mass deficits have been attributed to both increased marine-terminating outlet glacier discharge and to a more negative surface mass balance (SMB), primarily resulting from increased surface melting relative to accumulation (Rignot et al., 2008, 2011; van den Broeke et al., 2009; Zwally et al., 2011). The relative

contribution of each of these two components varies across the Arctic, but is presently approximately equal on the GIS (van den Broeke et al., 2009). A number of potential controls on marine-terminating outlet glacier behaviour have been identified (Figure 3), which we broadly classify as: (1) glacier-specific factors, which relate to the glaciological, topographic and geological setting of the glacier; and (2) climatic/oceanic forcing, including air and ocean temperatures, sea ice and precipitation. Important glacier-specific factors include subglacial topography and geology, fjord bathymetry and topography, sedimentation at the grounding line and glacier velocity, size, surface slope and catchment area (Figure 3) (Alley, 1991; Joughin et al., 2008b; Meier and Post, 1987). Theory suggests that changes in marine-terminating outlet glacier dynamics can occur independently of climatic/oceanic forcing (e.g. Alley, 1991; Meier and Post, 1987) and the importance of glacier-specific factors, particularly subglacial topography, has been highlighted by recent studies (Joughin et al., 2010, 2012; Thomas et al., 2009). Despite their apparent significance, however, the influence of glacier-specific factors on Arctic marine-terminating glacier behaviour is poorly understood.

In contrast, concerns over anthropogenic climate change in the 1990s resulted in an increasing focus on climatic/oceanic forcing factors and recent work has emphasized the widespread and synchronous nature of dynamic changes in many regions, particularly southeastern Greenland (e.g. Howat et al., 2008a; Murray et al., 2010). Consequently, this paper focuses on the climatic/oceanic drivers of marine-terminating Arctic outlet glacier dynamics and discusses three primary controls: air temperatures, ocean temperatures and sea ice concentrations (Figure 3). It should be noted, however, that these forcing factors are not independent (Figure 3) and that interconnections between them may significantly influence outlet glacier behaviour, yet many of these relationships are



**Figure 3.** Illustration of the primary climatic/oceanic forcing factors (black CAPS) and glacier-specific controls (white CAPS) thought to influence marine-terminating Arctic outlet glacier behaviour and mass balance. The major processes (black italics) and potential feedback mechanisms (white italics) are included. The role of meltwater enhanced basal sliding is represented with a dashed line as its influence on multi-year glacier behaviour remains equivocal.

*Image source:* Global Land Cover Facility ([www.landcover.org](http://www.landcover.org)).

poorly understood. We aim to: (1) review and summarize recent developments relating to each of these climatic/oceanic forcing factors; (2) highlight key uncertainties surrounding marine-terminating Arctic outlet glacier response to climatic/oceanic forcing; and (3) recommend directions for future research.

## II Arctic mass balance trends: 1990–2010

Rapid mass loss from Arctic masses has been documented since the early 1990s by numerous independent studies (Table 1) (e.g. Gardner et al., 2011; Krabill et al., 2004; Moholdt et al., 2010b; Rignot and Kanagaratnam, 2006; Velicogna and Wahr, 2006). Due to their remote

location and considerable size, mass balance is usually determined indirectly using remotely sensed data and/or SMB modelling. Considerable advances have been made in these techniques during the past 20 years, which have substantially improved our ability to quantify mass budgets and to assess the relative contribution of ice dynamics to mass loss (Krabill et al., 2004; Rignot and Kanagaratnam, 2006; van den Broeke et al., 2009; Velicogna and Wahr, 2006). At present, the primary techniques include Gravity Recovery and Climate Experiment (GRACE) data (e.g. Arendt et al., 2008; Bergmann et al., 2012; Jacob et al., 2012; Khan et al., 2010; Luthcke et al., 2006; Mémin et al., 2011; Velicogna, 2009; Velicogna and Wahr, 2006; Wouters et al., 2008), comparison

**Table 1.** Recent mass losses from the major glaciated regions and subregions of the Arctic. Data are first ordered according to regional mass loss rates and then according to mass loss rates from each subregion. The most recent estimates of total mass loss were used for each region and the latest values obtained from GRACE and SMB/D are presented for the GIS. Abbreviations are as follows: SMB – Surface mass balance; D – Discharge; GRACE – Gravity Recovery and Climate Experiment; SPOT – Système Pour l’Observation de la Terre; GNSS – Global Navigation Satellite System; RES – Radio Echo Sounding. \*Mass loss rates converted from  $Gt\ a^{-1}$  to  $km^3\ a^{-1}$ , assuming an ice density of  $0.917\ kg\ km^{-3}$  (IPCC, 2007).<sup>#</sup>This value includes peripheral ice caps and glaciers (Jacob et al., 2012).

Region	Subregion	Rate of mass loss ( $km^3\ a^{-1}$ )	Measurement period	Measurement method	Source
Greenland	Greenland Ice Sheet	$224.76 \pm 19^*$	1992–2009	SMB /D	Rignot et al., 2011
Greenland	Greenland Ice Sheet	$203.57 \pm 8.25^{*\#}$	2003–2010	GRACE	Jacob et al., 2012
Canadian Arctic	Ellesmere, Devon, Axel Heiberg and Baffin islands	$56.24 \pm 6.42^*$	2004–2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Canadian Arctic	Ellesmere, Devon, Axel Heiberg islands	$34.23 \pm 4.56^*$	2004–2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Canadian Arctic	Baffin Island	$22.0 \pm 4.28^*$	2004–2009	SMB/D, ICESat laser altimetry and GRACE	Gardner et al., 2011
Russian Arctic	Novaya Zemlya	$3.67 \pm 2$	2003–2010	GRACE	Jacob et al., 2012
Russian Arctic	Severnaya Zemlya	$0.92 \pm 2$	2003–2010	GRACE	Jacob et al., 2012
Russian Arctic	Franz Josef Land	$0 \pm 2$	2003–2010	GRACE	Jacob et al., 2012
Svalbard	Spitzbergen	$3.59 \pm 1.17$	2003–2008	ICESat laser altimetry and SPOT HRS 5 stereoscopic images	Moholdt et al., 2010b
Svalbard	Austfonna Ice Cap	$1.3 \pm 0.5$	2002–2008	ICESat laser altimetry, airborne laser altimetry, GNSS surface profiles and RES	Moholdt et al., 2010a
Svalbard	Barentsoya and Edgeoya	$0.46 \pm 0.30$	2003–2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b
Svalbard	Vestfonna Ice Cap	$0.39 \pm 0.20$	2003–2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b
Svalbard	Kvitoyjokeln ice cap	$0.32 \pm 0.08$	2003–2008	ICESat laser altimetry and topographic maps	Moholdt et al., 2010b

of SMB with outlet glacier discharge (Rignot et al., 2008, 2011; Rignot and Kanagaratnam, 2006; van den Broeke et al., 2009) and repeat laser or radar altimetry measurements (Abdalati et al., 2001; Krabill et al., 2004; Pritchard et al., 2009; Thomas et al., 2006, 2009).

The negative mass balance of the GIS has received particular attention and has been estimated via a number of techniques and for a range of time periods. The most recent values from GRACE (Jacob et al., 2012) and from the comparison of SMB with outlet glacier discharge (Rignot et al., 2011) are presented in Table 1. An important new trend is the rapid mass loss from the Canadian Arctic between 2007 and 2009, which made the archipelago the primary cryospheric contributor to eustatic sea level rise outside of the Greenland and Antarctic ice sheets (Table 1) (Gardner et al., 2011). Furthermore, the area has been highlighted as the largest potential contributor to ice loss and sea level rise of any glaciated region during the 21st century (Radić and Hock, 2011). Negative mass balance trends have also been documented in Svalbard (Hagen et al., 2009; Moholdt et al., 2010b; Nuth et al., 2010) and the Russian Arctic (Kotlyakov et al., 2010; Sharov et al., 2009) (Table 1). However, the mass balance of the Russian Arctic archipelagos have been comparatively poorly documented (Bassford et al., 2006). This represents a significant limitation to our understanding of the Arctic cryosphere, and highlights the need for further research in the region, as NZ, SZ and FJL account for approximately 20% of the glaciated area of the Arctic, excluding the GIS (Dowdeswell et al., 1997).

### *1 Spatial trends in Arctic mass balance*

Arctic mass balance trends have been spatially non-uniform, with many areas exhibiting slight growth at high elevations and rapid marginal thinning (e.g. Hagen et al., 2009; Pritchard et al., 2009; Sharov, 2010; Sharov et al., 2009;

Thomas et al., 2006, 2008; Zwally et al., 2011). Substantial thickening has been observed at high elevations on the GIS (Ettema et al., 2009; Johannessen et al., 2005; Thomas et al., 2006; Zwally et al., 2005); Austfonna ice cap, Svalbard (Bamber et al., 2004; Moholdt et al., 2010a, 2010b; Raper et al., 2005); the northern ice cap, NZ (Sharov et al., 2009); Tyndall and Windy ice domes in FJL; Schmidt and Vavilov ice caps in SZ (Sharov, 2010); and some Canadian Arctic ice caps (Abdalati et al., 2004; Mair et al., 2009). A number of potential explanations have been proposed for this interior thickening, including increased precipitation (Thomas et al., 2006; Zwally et al., 2005), possibly related to changes in sea ice extent (Bamber et al., 2004; Mair et al., 2009; Raper et al., 2005), long-term accumulation trends (Koerner, 2005; Moholdt et al., 2010a) and/or surge dynamics (Bevan et al., 2007). However, interior gains have been far outweighed by low-elevation thinning and marginal retreat (e.g. van den Broeke et al., 2009; Zwally et al., 2011), resulting in an overall negative mass balance in many regions (Table 1).

### *2 Dynamic contribution of marine-terminating outlet glaciers to mass loss*

In addition to rapid marginal thinning, peak losses have occurred on marine-terminating outlet glaciers (Moon and Joughin, 2008; Pritchard et al., 2009; Sole et al., 2008). On many of these glaciers, thinning rates of 10s of  $\text{m a}^{-1}$  have far exceeded surface melt rates, suggesting that thinning is largely 'dynamic' (i.e. resulting from changes in ice flow, rather than increased surface melting) (e.g. Abdalati et al., 2001; Burgess and Sharp, 2008; Krabill et al., 2004; Thomas et al., 2009). The contribution of glacier dynamics to recent mass deficits has been further emphasized by rapid retreat rates, which have reached kilometres per year on the GIS (e.g. Howat et al., 2008a; Joughin et al., 2008b, 2010; Moon and Joughin, 2008) and hundreds of metres per year elsewhere (e.g. Blaszczyk et al., 2009; Burgess



and Sharp, 2004; Nuth et al., 2010; Sharov, 2005). Furthermore, recent research has underscored the contribution of dynamic changes to decadal-scale losses, as initial perturbations at the glacier terminus may be rapidly transmitted to inland areas, producing widespread, substantial thinning (Howat et al., 2005, 2008b; Pritchard et al., 2009; Thomas et al., 2011; Zwally et al., 2011). This longer-term component of dynamic loss is an important emerging area of research and has the potential to be the primary component of the GIS contribution to 21st-century sea level rise (Price et al., 2011; Vieli and Nick, 2011).

Although the dynamics of marine-terminating outlet glaciers are now recognized as a key component of Arctic ice mass loss, they have also been highlighted as a principal area of uncertainty (IPCC, 2007). Specifically, the primary climatic/oceanic controls and the mechanisms by which they induce a dynamic response are yet to be fully understood (Howat et al., 2010; Sole et al., 2008; Vieli and Nick, 2011). The following sections review the three main climatic/oceanic controls identified to date, namely surface air temperatures, ocean temperatures and sea ice concentrations, and discuss the primary linkages between these factors (Figure 3). All three forcing factors have undergone marked changes in recent years, which have been linked to both recent climatic warming (ACIA, 2004; IPCC, 2007) and to the onset of a negative phase of the North Atlantic Oscillation (NAO) in the mid-1990s (e.g. Stern and Heide-Jørgensen, 2003; Gerdes et al., 2003; Hurrell et al., 2003; Holliday et al., 2008).

### III Air temperature forcing


Arctic air temperatures have risen substantially since the mid-1990s (ACIA, 2004; Hanna et al., 2008; IPCC, 2007), although they are not unprecedented at decadal timescales (Box et al., 2009; Chylek et al., 2006). We present a new synthesis of air temperature data to investigate the spatial

distribution of Arctic warming between 1990 and 2010 and to visualize this trend in terms of both magnitude and statistical significance (Figure 4). Linear trends were calculated from annual air temperature series, which were compiled from meteorological station data of varying temporal resolution (three-hourly to monthly). In order to account for missing values, three-hourly data were used only if: (1) no more than two consecutive records were missing in a day; and (2) no more than three records in total were missing in a day. Daily data were only used if values were available for 22 or more days per month and monthly values were used only if data were available for all months of the year (Cappelen, 2011).

Results suggest that warming has been greatest at coastal stations surrounding Baffin Bay and the Davis Strait (Figure 4), which is consistent with dramatic mass loss from the Canadian Arctic between 2004 and 2009 (Gardner et al., 2011). Significant warming has also occurred in the Kara Sea region, particularly on FJL (Figure 4), although data coverage is comparatively sparse. Warming from the mid-1990s has been linked to negative SMB on a number of Arctic ice masses, particularly the GIS (e.g. Abdalati and Steffen, 2001; Bhattacharya et al., 2009; Box et al., 2006; Ettema et al., 2009; Hanna et al., 2008; Mote, 2007). However, while warming directly affects SMB, a key recent development has been to consider the potential impact of meltwater on outlet glacier dynamics.

#### *I Air temperatures, meltwater production and ice velocities on temperate and polythermal glaciers*

The relationship between air temperatures, meltwater supply and ice velocities has been well documented on temperate glaciers (e.g. Fountain and Walder, 1998; Iken and Bindschadler, 1986; Willis, 1995), but had not been extensively considered on large Arctic ice masses until relatively recently. On temperate glaciers, surface meltwater is thought to access large portions of the




**Figure 4.** Linear trend in mean annual air temperatures between 1990 and 2010 for selected Arctic meteorological stations. Symbol colour shows the magnitude of the linear trend in °C per year between 1990 and 2010. Symbol size shows the  $R^2$  value of the relationship: a larger symbol represents a larger  $R^2$  value and therefore the trend line better fits the data. Meteorological stations were selected according to data availability for the study period.

*Meteorological data sources:* Danish Meteorological Institute, weather and climate data from Greenland 1958–2010; Norwegian Meteorological Institute, Klima climate database; Royal Netherlands Meteorological Institute, Climate Explorer; Scientific Research Institute of Hydrometeorological Information, World Data Center – Baseline Climatological Data Sets; and National Climate Data and Information Archive, Canadian Daily Climate Data.

glacier bed during the melt season, resulting in elevated basal water pressures, reduced basal drag and enhanced ice motion (e.g. Fountain and Walder, 1998; Iken and Bindenschadler, 1986; Kamb, 1987; Nienow et al., 1998; Willis, 1995). As the melt season progresses, continued meltwater input promotes the development of a more efficient subglacial drainage system, which lowers basal water pressures and reduces the sensitivity of glacier velocities to additional melt (Figure 5) (e.g. Nienow et al., 1998; Willis,

1995). Recent studies have demonstrated a similar relationship on polythermal glaciers in the Canadian Arctic (e.g. Bingham et al., 2003, 2008; Boon and Sharp, 2003; Copland et al., 2003) and in Svalbard (Nuttall and Hodgkins, 2005; Rippin et al., 2005; Vieli et al., 2004). In particular, extensive investigations on John Evans Glacier (JEG), Ellesmere Island, Canada, showed that surface meltwater could rapidly access the bed through predominantly cold ice and cause substantial seasonal acceleration



**Figure 5.** Idealized seasonal evolution of glacier response to meltwater inputs. The graph illustrates the theoretical response of outlet glacier velocities to meltwater inputs during the melt season. The bottom panels illustrate an idealized plan view of the subglacial hydrological system at different stages of the melt season (bottom panels modified from Fountain and Walder, 1998). Individual glacier response to meltwater forcing may vary significantly from this idealized situation.

(Copland et al., 2003; Bingham et al., 2003, 2005, 2008).

## 2 Surface meltwater and ice velocities in the GIS ablation zone

Until a decade ago, it was largely assumed that penetration of surface meltwater to the bed of

large Arctic ice masses would be minimal and that its effect on ice velocities would be limited, especially on the GIS (Copland et al., 2003; Hodgkins, 1997; Zwally et al., 2002). This viewpoint was radically altered by GPS measurements from Swiss Camp in the west Greenland ablation zone, which first demonstrated a close correspondence between surface meltwater inputs and ice

velocities (Zwally et al., 2002). Here we define the ablation zone as areas that experience melt, with the exception of fast-flowing, marine-terminating outlet glaciers, which are discussed separately (section III.3), due to their differing response to meltwater inputs. Results from Swiss Camp showed that velocities closely followed seasonal and interannual variations in surface meltwater production, as previously observed on temperate glaciers, and this was attributed to meltwater-enhanced basal sliding (Zwally et al., 2002). Most importantly, the study highlighted meltwater-enhanced basal lubrication as a potential mechanism for rapid, dynamic and widespread response of the GIS to atmospheric warming (Zwally et al., 2002).

The work of Zwally et al. (2002) was supported by subsequent results from the west Greenland ablation zone, which provided further evidence of rapid coupling between seasonal meltwater inputs and ice velocities (e.g. Bartholomew et al., 2010, 2011; Catania and Neumann, 2010; Das et al., 2008; Joughin et al., 2008a; van de Wal et al., 2008). Studies also identified supraglacial lake drainage events as a potential mechanism for rapid transfer of meltwater to the bed (e.g. Das et al., 2008; Krawczynski et al., 2009). Large volumes of water released during drainage events may promote crevasse propagation through the full ice thickness by offsetting rapid refreezing and maintaining high water pressures at the crevasse tip (Alley et al., 2005; Krawczynski et al., 2009; van der Veen, 1998, 2007). Drainage events have immediately preceded velocity increases in the west Greenland ablation zone (Box and Ski, 2007; Das et al., 2008; McMillan et al., 2007), on land-terminating west Greenland outlet glaciers (Shepherd et al., 2009; Sneed and Hamilton, 2007) and on JEG (Bingham et al., 2003; Boon and Sharp, 2003; Copland et al., 2003), providing empirical support for their role in meltwater delivery to the bed.

The potential impact of surface meltwater inputs on the GIS was also explored using


numerical modelling, which predicted far greater losses with enhanced basal sliding (Huybrechts and de Wolde, 1999; Parisek and Alley, 2004; van de Wal and Oerlemans, 1997). This occurred via a number of proposed feedback mechanisms, which are illustrated for an idealized section of the GIS (Figure 6). Specifically, feedbacks could develop between glacier acceleration, dynamic thinning and surface melting: increased basal sliding would promote dynamic thinning and bring a greater portion of the ice sheet into the ablation zone, thus exposing a greater area to melting and enhanced lubrication (Figure 6) (Parisek and Alley, 2004).

### *3 Surface meltwater and marine-terminating Arctic outlet glacier dynamics*

The close coupling between surface meltwater and ice velocities observed in the GIS ablation zone led to increased consideration of the influence of meltwater on marine-terminating outlet glacier dynamics (e.g. Hall et al., 2008; Krabill et al., 2004). This was further motivated by the concurrence of the onset of marine-terminating Arctic glacier retreat from the mid-1990s with atmospheric warming (e.g. Bevan et al., 2012a; Dyurgerov and McCabe, 2006; Howat and Eddy, 2011) and the coincidence of substantial changes in glacier dynamics with elevated air temperatures (e.g. Howat et al., 2008a; Moon and Joughin, 2008; Rignot and Kanagaratnam, 2006).

Recent results from marine-terminating Arctic outlet glaciers appear to support meltwater-enhanced basal lubrication as a mechanism for ice acceleration at subannual timescales: glacier velocities in the Uummannaq region of west Greenland (Howat et al., 2010) and on Duvebeen, Austfonna (Figure 1) (Dunse et al., 2012), closely corresponded to the seasonal melt cycle. Similarly, results from Petermann Glacier (Figures 1 and 2) (Nick et al., 2012) and Dagaard Jensen Gletscher (Figure 1) (Bevan et al., 2012b) suggest that seasonal velocities primarily reflect variations in surface meltwater availability, and





**Figure 6.** Proposed feedback mechanisms between surface meltwater availability, basal sliding and ice sheet geometry for an idealized section of the GIS. Atmospheric warming may increase surface meltwater input to the bed, resulting in enhanced basal sliding and transfer of a greater portion of the outlet glacier to the ablation zone. Further feedbacks may then develop between dynamic thinning, inland migration of basal sliding and ice acceleration. The response of individual sections of the ice sheet may vary significantly from these idealized theoretical responses.

data from Helheim Glacier (HH) (Figure 1) indicate that surface meltwater can be transmitted to the bed within 12–36 hours (Andersen et al., 2010a).

Despite an apparent relationship at seasonal or shorter timescales, however, the influence of meltwater-enhanced basal lubrication on interannual marine-terminating outlet glacier behaviour remains equivocal (e.g. Bingham et al., 2003; McFadden et al., 2011; Seale et al., 2011; van de Wal et al., 2008; Vieli et al., 2004). Evidence from the GIS suggests that meltwater input to the bed may have a limited impact on interannual velocity changes on fast-flowing marine-terminating outlet glaciers and that ice flow may be more responsive to conditions at the

ice-ocean interface (Joughin et al., 2008a; Nick et al., 2009). A similar pattern has been observed on JEG (Bingham et al., 2003) and Hansbreen, Spitzbergen (Figure 1) (Viellet et al., 2004), where periods of high melt coincided with reduced seasonal acceleration or even deceleration. Furthermore, numerical modelling results from HH (Nick et al., 2009) suggest that changes in frontal position, as opposed to meltwater-enhanced basal lubrication, are the dominant control on interannual behaviour. Thus, evidence suggests that meltwater-enhanced basal lubrication may significantly influence marine-terminating outlet glacier dynamics at subannual timescales, but its role in driving interannual retreat remains uncertain.

To date, research into the influence of meltwater on marine-terminating outlet glacier dynamics has predominantly focused on enhanced basal lubrication. However, meltwater may also influence dynamics by promoting crevasse propagation at the terminus and/or lateral margins (Figure 3), which together could reduce resistive stresses and promote glacier retreat (Andersen et al., 2010b; Sohn et al., 1998; van der Veen, 1998; van der Veen et al., 2011; Vieli et al., 2007). This partly agrees with model results from JI, which suggest that increased crevasse water levels can partially reproduce observed patterns of retreat and acceleration, but this may also reflect the choice of calving model (Vieli and Nick, 2011). Numerical modelling studies also suggest that acceleration at Jakobshavn Isbrae (JI), west Greenland, may have resulted from weakening at its lateral margins, potentially due to hydrofracturing and/or meltwater-induced warming of the ice (van der Veen et al., 2011). Thus, while the role of meltwater-enhanced fracture as a primary trigger of retreat remains equivocal, this mechanism warrants further consideration given the sensitivity of marine-terminating glaciers to changes at the terminus (Nick et al., 2009; Vieli and Nick, 2011).

#### *4 Subglacial drainage systems of large Arctic ice masses*

Research into the subglacial hydrology of Arctic ice masses has predominantly focused on land-terminating sections, but recent advances, particularly from the GIS, may provide insight into the comparative insensitivity of marine-terminating outlet glaciers to meltwater-enhanced basal lubrication at interannual timescales. Although the subglacial hydrology of marine-terminating outlet glaciers is comparatively poorly understood and the response of individual glaciers may vary significantly, observations suggest that the seasonal evolution of the subglacial drainage system is very similar to that observed on temperate,

polythermal and land-terminating outlet glaciers and sections of the GIS ablation zone: the subglacial drainage system is thought to evolve during the melt season, causing variation in the sensitivity of ice velocities to meltwater inputs (Figure 5) (e.g. Bartholomew et al., 2010, 2011; Copland et al., 2003; Dunse et al., 2012; Howat et al., 2010; Shepherd et al., 2009; Sole et al., 2011; Vieli et al., 2004). Early in the melt season, the drainage system may be relatively inefficient (Figure 5) (Bartholomew et al., 2010; Bingham et al., 2003; Kamb, 1987; Price et al., 2008). Consequently, meltwater can rapidly increase basal water pressures, causing rapid ice acceleration and surface uplift (Bartholomew et al., 2010; Bingham et al., 2005; Copland et al., 2003). As the melt season progresses, continued inflow of surface meltwater may promote the development of a more efficient, channellized drainage system which operates at lower basal water pressures (Figure 5) (Bingham et al., 2003, 2006; Kamb, 1987; Palmer et al., 2011; Shepherd et al., 2009; Sole et al., 2011). Thus, the sensitivity of ice velocities to surface melt may decline and only large meltwater inputs may induce substantial velocity change (Figure 5) (Bartholomew et al., 2010; Dunse et al., 2012; Schoof, 2010; Shepherd et al., 2009). The primary implication of these results is that ice velocities depend not only on surface meltwater inputs, but also on the subglacial hydrological system.

The evolution of the subglacial drainage system has important implications for the response of marine-terminating outlet glaciers to interannual variations in meltwater availability and atmospheric warming (Price et al., 2008; Schoof, 2010; Sundal et al., 2011; van de Wal et al., 2008). As observed at seasonal timescales, continually high meltwater inputs are likely to promote the formation of an efficient basal drainage system, operating at low water pressures (Figure 5). Consequently, increased meltwater input at interannual timescales may not necessarily equate to increased ice velocities, and may even cause deceleration above critical thresholds of water

supply (Schoof, 2010; Sundal et al., 2011; Vieli et al., 2004). This is consistent with empirical results from Kangiata Nunata Sermia, southwestern Greenland, where meltwater-induced summer speed-up events are thought to contribute little to annual ice velocities, partly because they are offset by the deceleration associated with the formation of an efficient subglacial system (Sole et al., 2011). The key conclusion of these findings is that the evolution of the hydrological system may act as a buffer against accelerated ice loss through meltwater-enhanced basal sliding in response to increased melt and atmospheric warming (Price et al., 2008; Schoof, 2010; Vieli et al., 2004).

## IV Oceanic forcing


While atmospheric warming has received substantial scientific attention, oceanic forcing has been recently recognized as a key control on marine-terminating outlet glacier dynamics. This was partly instigated by results from the GIS (e.g. Moon and Joughin, 2008; Pritchard et al., 2009; Sole et al., 2008), where retreat rates were approximately two orders of magnitude greater on marine-terminating glaciers (10s to 1000s of  $\text{m a}^{-1}$ ) than on their land-terminating counterparts (0.1 to 1  $\text{m a}^{-1}$ ) (Figure 7). A similar pattern has been observed elsewhere in the Arctic, including Austfonna ice cap (Dowdeswell et al., 2008), Devon Ice Cap (Burgess and Sharp, 2004, 2008; Dowdeswell et al., 2004; Shepherd et al., 2007) and in Arctic Alaska (Arendt et al., 2006). Furthermore, thinning rates have been greatest on glaciers occupying deep bedrock troughs (Thomas et al., 2009), which may allow warm, subsurface Atlantic Water (AW) from the continental shelf to access the glacier termini (e.g. Rignot et al., 2010; Straneo et al., 2010, 2011). Oceanic forcing may be of particular concern in the near future, as model predictions suggest that ocean temperatures around the GIS may warm by 1.7–2°C by 2100 (Yin et al., 2012).

### *1 Submarine melting at marine-terminating outlet glacier termini*

Measurements of submarine melt rates at the termini of marine-terminating glaciers are rare, but estimates suggest that rates range between  $0.7 \pm 0.2$  and  $3.9 \pm 0.8$  m per day in central west Greenland (Rignot et al., 2010) and  $4.34 \pm 0.94$  m per day at JI (Motyka et al., 2011). Substantially higher melt rates of 6.9 to 12.4 m per day have been estimated at LeConte Glacier, Alaska (Figure 1) (Motyka et al., 2003), probably reflecting its comparatively southerly location. These results highlight the potential sensitivity of marine-terminating glaciers to oceanic warming, which could influence outlet glacier dynamics via a number of mechanisms (Figure 8). First, enhanced submarine melting may cause grounding-line retreat at floating and grounded margins, potentially resulting in further ungrounding and the development of positive feedbacks if retreat occurs into deeper water (Howat et al., 2008a; Joughin et al., 2008b; Meier and Post, 1987; Nick et al., 2012; Vieli and Nick, 2011; Vieli et al., 2001). Second, oceanic warming may cause rapid thinning of floating termini (e.g. Motyka et al., 2011; Nick et al., 2012; Thomas, 2004) and the formation of deeply incised basal channels (Rignot and Steffen, 2008), which together make the termini more vulnerable to full thickness fracture and eventual disintegration (Figure 8). Third, submarine melting may influence the terminus geometry and calving rates by undercutting at the grounding line and/or waterline (Figure 8) (Benn et al., 2007; Vieli et al., 2002).

### *2 Oceanic controls on marine-terminating glacier dynamics*


Our understanding of oceanic forcing has been largely developed from observations from the GIS, where warming has immediately preceded the retreat and acceleration of a number of



**Figure 7.** Mean rate of Greenland outlet glacier frontal position change ( $\text{m a}^{-1}$ ) grouped according to terminus type. The mean rate of retreat, advance and net frontal position change were calculated for land-terminating and tidewater glacier termini and are shown in the bars above. Values were calculated for three time periods (1992–2000, 2000–2006 and 2006–2007) and maximum rates of retreat/advance are given in brackets above the corresponding bar. Mean values are calculated from a sample of 139 (1992–2000), 169 (2000–2006) and 154 (2006–2007) tidewater glaciers, and 10 (1992–2000), 14 (2000–2006) and 13 (2006–2007) land-terminating glaciers. Glaciers terminating in ice shelves were excluded from the analysis, as data were only available from three glaciers for 1992–2000 and 2000–2006 and no data were available for 2006–2007.

Source: Data provided by T. Moon, 2011 (Moon and Joughin, 2008).





**Figure 8.** Illustration of the influence of oceanic warming and submarine melting on outlet glacier dynamics and geometry for (A) an initially floating terminus and (B) an initially grounded terminus. In (A), feedbacks may develop between submarine melting, grounding-line retreat, thinning and calving front retreat. In (B), changes in terminus geometry may initiate feedbacks between grounding-line/terminus retreat, thinning and floatation.

marine-terminating outlet glaciers (e.g. Bevan et al., 2012a; Hanna et al., 2009; Holland et al., 2008; Motyka et al., 2011; Murray et al., 2010; Rignot et al., 2012). This was first investigated in detail at JI, which was one of the earliest and most significant contributors to recent GIS mass

losses (Joughin et al., 2004, 2008c; Motyka et al., 2010, 2011; Rignot and Kanagaratnam, 2006; Thomas et al., 2003). Following 50 years of comparative stability (Csatho et al., 2008; Sohn et al., 1998), JI's floating terminus began to retreat in October 1998 (Luckman and Murray, 2005) and

subsequent periods of acceleration often coincided with the loss of sections of its tongue (Joughin et al., 2004, 2008c). Initial retreat was accompanied by rapid thinning, which may have ungrounded the tongue from its underlying pinning points, and caused a substantial reduction in resistive stresses (Joughin et al., 2004; Thomas, 2004; Thomas et al., 2003). This may have initiated feedbacks between retreat, dynamic thinning and acceleration, which led to the disintegration of the ice tongue by spring 2003 (Joughin et al., 2004, 2008c; Thomas, 2004).


The underlying driver(s) of mass losses at JI remain subject to debate, but evidence suggests that oceanic warming, rather than increased air temperatures, was the primary cause (Holland et al., 2008; Motyka et al., 2010, 2011; Thomas, 2004). Thinning rates on JI's floating tongue far exceeded estimated surface melt rates and closely followed substantial subsurface ocean warming, which is thought to have increased basal melt rates by 25% (Holland et al., 2008; Motyka et al., 2011; Thomas et al., 2003). Estimates suggest that the resultant thinning was sufficient to destabilize the ice tongue and to initiate rapid mass loss (Motyka et al., 2011). Numerical modelling results agree with these findings and suggest that increased submarine melting is capable of triggering the behaviour observed at JI, but that dynamic feedbacks are also required (Vieli and Nick, 2011).

Subsequent to retreat at JI, marine-terminating outlet glaciers in southeastern Greenland followed a similar progression of dynamic change (e.g. Howat et al., 2007, 2008a; Joughin et al., 2008b; Luckman et al., 2006). Losses began with retreat, thinning and acceleration proportional to retreat, which suggests that changes also resulted from a loss of resistive stresses at the terminus (Howat et al., 2005, 2007, 2008a). The trigger for these changes remains equivocal, with both air temperatures (Box et al., 2009; Hanna et al., 2008) and ocean temperatures (Hanna et al., 2009; Murray et al., 2010; Seale et al., 2011) increasing substantially prior to retreat.

However, the initiation of glacier response at the terminus (Howat et al., 2005, 2007, 2008a) suggests that meltwater-enhanced basal lubrication was unlikely to be the primary trigger and that forcing factors operating at the calving front, such as oceanic warming, were the more likely cause. This is consistent with numerical modelling results from HH, which suggested that inter-annual glacier dynamics are comparatively insensitive to enhanced basal lubrication, but are acutely sensitive to calving front perturbations (Nick et al., 2009).

### *3 Marine-terminating outlet glacier dynamics and Atlantic Water distribution*

An important emerging theme has been the relationship between marine-terminating outlet glacier dynamics and variations in the distribution and properties of warm Atlantic Water (AW) (Andersen et al., 2012; Holland et al., 2008; Murray et al., 2010; Straneo et al., 2010, 2011). Until recently, it was assumed that oceanic changes at the continental shelf could be transmitted into outlet glacier fjords, but this was largely untested (Mortensen et al., 2011; Straneo et al., 2010). However, recent studies have shown that AW can access the fjords of a number of large outlet glaciers in Greenland (Christoffersen et al., 2011; Holland et al., 2008; Johnson et al., 2011; Mayer et al., 2000; Straneo et al., 2010, 2011) and Svalbard (Nilsen et al., 2008). These results marked a significant advance in our understanding, as they demonstrated that rapid connections could exist between marine-terminating outlet glaciers and oceanic variability in the northern North Atlantic, particularly via deep fjords (Straneo et al., 2010). This conclusion was supported by the coincidence of glacier retreat in southeastern Greenland in the early 2000s with AW incursion onto the coast (Christoffersen et al., 2011; Murray et al., 2010; Seale et al., 2011) and provides a plausible mechanism for widespread and synchronous retreat.



**Figure 9.** Schematic illustrating the circulation pattern and water properties within a large Arctic outlet glacier fjord. Fjord circulation and water mass depths are based on conditions within Helheim Glacier fjord (Straneo et al., 2011). The primary controls on fjord circulation are thought to be water properties at the continental shelf, wind/storm tracks and glacial meltwater input.

#### 4 Marine-terminating outlet glacier dynamics and fjord circulation

Recent research into the role of AW has led to increased consideration of the factors controlling its distribution within glacial fjords. A number of possible controls have been identified (Figure 9), including: the temperature, salinity and volume of subtropical waters at the continental shelf; along-shore wind patterns; storm tracks; and fjord stratification (Christoffersen et al., 2011; Nilsen et al., 2008; Straneo et al., 2010, 2011). Fjord circulation can also be influenced by subglacial meltwater, which forms a rising plume of cool, buoyant water at the calving front and promotes a compensatory inflow of warmer water at depth (Figure 9) (Motyka et al., 2003, 2011; Straneo et al., 2011). Thus, plumes may substantially increase submarine melt rates (Jenkins, 2011; Motyka et al., 2003; Seale et al., 2011) and model results suggest that melt increases linearly with oceanic warming and to the power of one-third with subglacial discharge (Jenkins, 2011; Xu et al., 2012). A key implication of this relationship is

that positive feedbacks could develop, whereby atmospheric warming increases subglacial discharge and ice sheet runoff, which strengthens the plume and enhances submarine melt rates (Seale et al., 2011). Feedbacks between glacier runoff and ocean properties have been identified as a potential trigger for recent retreat in southeastern Greenland (Murray et al., 2010; Seale et al., 2011) and variations in meltwater production may be an important control on AW distribution in the region (Murray et al., 2010).

#### V Sea ice forcing

The increasing focus on oceanic forcing has led to further consideration of the influence of sea ice on marine-terminating Arctic outlet glacier behaviour (Figure 3). Although sea ice is discussed separately, it should be noted that it is influenced by both air and ocean temperatures (Figure 3) and that these factors are not independent. It should also be noted that sea ice concentrations may significantly affect SMB, through their influence on accumulation and ablation patterns (Figure 3) (e.g. Bamber et al., 2004;

Rennermalm et al., 2009). The influence of sea ice on marine-terminating Arctic outlet glacier dynamics was first documented in northern Greenland, where semi-permanent fast-ice contributed significantly to the stability of several marine-terminating outlet glaciers (Higgins, 1989, 1990; Mayer et al., 2000; Reeh et al., 2001; Weidick, 1975). Fast-ice was thought to promote glacier stability by suppressing calving and by preventing calved material from moving away from the terminus (Higgins, 1990; Reeh et al., 2001). In contrast, periods of fast-ice disintegration were accompanied by rapid calving and release of trapped ice. Early investigations suggested that fast-ice break-up occurred at decadal intervals, when summer temperatures were exceptionally warm (Higgins, 1989, 1990; Reeh et al., 2001), but this pattern has changed substantially in recent years, with disintegration now occurring several times per decade (Hughes et al., 2011).

### *1 Sea ice influence on the seasonal calving cycle*


Recent studies have investigated the influence of sea ice on calving rates at more southerly Greenland glaciers (Ahn and Box, 2010; Howat et al., 2010), particularly on JI (Amundson et al., 2010; Joughin et al., 2008c; Sohn et al., 1998). As in northern Greenland, sea ice concentrations at JI appear to influence the timing and nature of calving events, but this occurs on seasonal, as opposed to decadal, timescales (Amundson et al., 2010; Joughin et al., 2008c). In winter, sea ice binds together icebergs to form a semi-rigid, seasonal ice shelf, or *mélange*, which is pushed along the fjord as a coherent mass by the advancing calving front (Figure 10) (Amundson et al., 2010). The *mélange* suppresses calving rates by up to a factor of six and alters the terminus geometry and near-front stress fields, causing seasonal terminus advance and deceleration (Amundson et al., 2010; Joughin et al., 2008c; Sohn et al., 1998). Conversely, springtime

*mélange* disintegration allows high rates of summer calving to commence, which initiates seasonal retreat and acceleration (Figure 10) (Ahn and Box, 2010; Amundson et al., 2010; Howat et al., 2010; Joughin et al., 2008c). A similar relationship has been documented on the Agassiz Ice Cap, Ellesmere Island, Arctic Canada, where peak glacier velocities have coincided with seasonal sea ice disintegration (Williamson et al., 2008). However, observations also indicated that sea ice weakening and/or thinning, as opposed to complete disintegration, may be sufficient to initiate seasonal acceleration (Williamson et al., 2008).

### *2 Sea ice influence on interannual marine-terminating outlet glacier behaviour*

Observations from JI have contributed substantially to our understanding of sea ice forcing at seasonal timescales, but have also highlighted its potential influence on interannual behaviour of marine-terminating outlet glaciers (Joughin et al., 2008c). Initial retreat at JI began within one year of the onset of sea ice decline in the surrounding Disko Bay (Joughin et al., 2008c). Estimates suggest that the extension of ice-free conditions by one or two months may have been sufficient to trigger the initial retreat by extending the duration of seasonally high calving rates (Joughin et al., 2008c). This is consistent with numerical modelling results which demonstrated that reduced *mélange* duration could trigger rapid retreat at JI, although it could not replicate the magnitude of subsequent seasonal variations in terminus position (Vieli and Nick, 2011). A similar response has been observed in the Uummannaq region (Howat et al., 2010) and at KG (Christoffersen et al., 2011; Seale et al., 2011), where interannual retreats also followed sea ice decline. It is thought that delayed winter sea ice formation at KG (Christoffersen et al., 2011; Seale et al., 2011) and early *mélange* clearance in the Uummannaq region (Howat et al., 2010) may have






**Figure 10.** Illustration of the influence of sea ice and mélangé formation on Arctic outlet glacier dynamics during (A) mélangé formation at the end of the calving season and (B) mélangé disintegration at the start of the calving season. In (A) the mélangé binds together material within the fjord, thus suppressing calving and promoting seasonal advance. In (B) mélangé disintegration allows seasonally high calving rates to commence and promotes glacier retreat.

initiated glacier retreat by extending the calving season.

Although the influence of sea ice on marine-terminating outlet glacier behaviour has been little studied outside of the GIS, Arctic sea ice has declined markedly in recent years (e.g.

Kwok and Rothcock, 2009; Rodrigues, 2009; Serreze et al., 2009) and its influence may become increasingly widespread if current losses continue. On the basis of the relationships observed in Greenland, we suggest that sea ice decline may affect glacier dynamics via two



**Figure 11.** Multi-model mean sea ice concentration (%) for January to March (JFM) and June to September (JAS) in the Arctic for the periods (a) 1980–2000 and (b) 2080–2100 for the SRES A1B scenario. The dashed white line indicates the present-day 15% average sea ice concentration limit. Note the substantial reduction in summer sea ice concentrations predicted across the Arctic by 2100, which may extend seasonally ice-free conditions in southerly areas and may result in a transition from multi-year fast-ice to seasonal sea ice disintegration in northern regions.

Source: Modified from IPCC (2007) and Flato et al. (2004).

potential mechanisms: (1) seasonal calving may be extended in areas which currently experience seasonally ice-free conditions; and (2) areas currently characterized by interannual fast-ice may transition to a seasonal sea ice loss. We suggest that the former process may become increasingly significant on the eastern and central-western Greenland coast, on the western coasts of NZ and Svalbard and in the southern Canadian Arctic, where the ice-free season has extended markedly during the past 30 years (Rodrigues, 2008) and losses are predicted to continue during the 21st century (Figure 11) (ACIA, 2004; IPCC, 2007). This mechanism may eventually cease, however, if areas become perennially ice-free. The latter process may become increasingly important on the coasts of northeastern Greenland, northeastern Svalbard, eastern NZ, southern FJL and the northern Canadian Arctic, where sea ice concentrations are predicted to decline markedly by 2100 (Figure 11) (ACIA, 2004; IPCC, 2007). Observations suggest that this may already be occurring in northeastern Greenland, where fast-ice break up has occurred several times in the past

decade (Hughes et al., 2011), in comparison to the decadal intervals recorded by earlier work (Higgins, 1989, 1990; Reeh et al., 2001).

## VI Key uncertainties and future directions for research

Despite recent advances, the response of marine-terminating outlet glaciers to climatic/oceanic forcing continues to be an area of rapidly developing research, and significant uncertainties remain over the relative importance of each forcing factor and the mechanisms by which these factors influence glacier dynamics (Howat et al., 2010; Sole et al., 2008; Vieli and Nick, 2011). The following subsections outline the primary uncertainties surrounding marine-terminating Arctic outlet glacier behaviour and highlight key areas for future research.

### *I Spatial variation in the relative importance of climatic/oceanic forcing factors*


Our understanding of marine-terminating Arctic outlet glacier response to climatic/oceanic

forcing has been primarily based on observations from a small number of Greenland outlet glaciers, with the majority of research focusing on JI and southeastern Greenland, particularly HH and KG. Consequently, it is uncertain whether the relationships observed at these locations can be extrapolated to other Arctic regions and/or whether recent changes represent a longer-term trend or shorter-term variability (Price et al., 2011; Vieli and Nick, 2011). Although glaciers within certain regions have shown some common response to climatic/oceanic forcing, most notably southeastern Greenland (Bjørk et al., 2012; Howat et al., 2008a; Murray et al., 2010), this pattern is far from ubiquitous. Results from west Greenland found no correlation between retreat and climatic/oceanic forcing for a sample of 59 marine-terminating outlet glaciers (McFadden et al., 2011) and comparison of 15 major Greenland outlet glaciers between 1985 and 2011 showed some common response to forcing, but also highlighted several notable differences (Bevan et al., 2012a). Furthermore, assessment of decadal and interannual velocity changes on >200 major Greenland outlet glaciers demonstrated substantial variations in glacier behaviour at both regional and local scales, and highlighted the importance of glacier-specific factors (Moon et al., 2012). In contrast to the GIS, observations in the Canadian Arctic (Gardner et al., 2011) and Novaya Zemlya (Moholdt et al., 2012) have found no difference between area-averaged thinning rates in land- and marine-terminating basins (Gardner et al., 2011). Moreover, the longer-term evolution of HH, KG and JI has differed markedly following their earlier mass losses (Howat et al., 2011; Thomas et al., 2011) and numerical modelling studies indicate that marine-terminating outlet glaciers can rapidly adjust to short-term calving front perturbations (Viel and Nick, 2011). Together, this evidence suggests that the relative importance of climatic/oceanic controls varies across the Arctic and that present theories of outlet glacier response to forcing cannot be universally applied to all glaciers, regions

or ice masses. We therefore draw attention to the danger of extrapolating recent rapid mass losses from a small number of glaciers and highlight the need for continued research into the climatic/oceanic drivers of marine-terminating outlet glacier behaviour on each of the major Arctic ice masses.

## 2 Glacier-specific factors

Results from the GIS have highlighted the substantial variation in marine-terminating outlet glacier response to climatic/oceanic forcing (McFadden et al., 2011; Moon et al., 2012), and the role of glacier-specific controls, particularly fjord geometry and basal topography, is being increasingly recognized (Bevan et al., 2012a; Howat and Eddy, 2011; Joughin et al., 2010, 2012; Nick et al., 2009; Thomas et al., 2009). Traditional theories of tidewater glacier dynamics and ice sheet instability suggest that a reverse basal slope may initiate rapid retreat via a series of positive feedbacks, as the glacier terminus retreats into progressively deeper water (Figure 12) (e.g. Hughes, 1986; Joughin et al., 2008b; Meier and Post, 1987; Vieli et al., 2001, 2002; Weertman, 1974). This behaviour may occur independently of climatic/oceanic forcing (e.g. Alley, 1991; Pfeffer, 2003), but may also be initiated by perturbations at the calving front (e.g. Howat et al., 2008a; Joughin et al., 2008b; Meier and Post, 1987; Nick et al., 2009; Pfeffer, 2007). However, the influence of overdeepenings on glacier dynamics remains subject to debate, and recent modelling results suggest that stable grounding-line positions can be achieved on a reverse bedrock slope (Gudmundsson et al., 2012; Nick et al., 2010). Furthermore, the importance of other glacier-specific factors, such as variations in fjord width, is being increasingly acknowledged (Jamieson et al., 2012). Assessing the role of glacier-specific controls is a key area for future study, as inadequate consideration of these factors may lead to substantial errors in estimates of glacier response to climatic/oceanic



**Figure 12.** Illustration of feedbacks between glacier retreat, dynamic thinning and ice acceleration during retreat into progressively deeper water. Initial retreat reduces resistive stresses acting on the outlet glacier, promoting dynamic thinning and terminus floatation, which in turn makes the terminus increasingly vulnerable to fracture and further retreat. Positive feedbacks may also develop between grounding-line retreat and submarine melt rates. These feedbacks may occur independently of climatic/oceanic forcing, but may also be triggered by forcing.

forcing and their contribution to sea level rise. A full analysis is, however, currently constrained by limited data availability.

### 3 Quantitative assessment of marine-terminating outlet glacier response to climatic/oceanic forcing

Even on comparatively well-studied sections of the GIS, previous studies have tended to infer causality from the coincidence of climatic/oceanic change and marine-terminating outlet glacier response (e.g. Luckman et al., 2006; Moon and Joughin, 2008). As a consequence, the mechanisms linking climatic/oceanic forcing and glacier dynamics are often poorly understood (Nick et al., 2009; Vieli and Nick, 2011) and the extent to which forcing can explain glacier behaviour has not been extensively assessed. This has been improved in recent years through the development of numerical models focusing on the response of individual outlet glaciers to forcing

(Nick et al., 2009; Vieli and Nick, 2011). However, marine-terminating outlet glacier dynamics are not yet adequately represented in ice sheet scale models (Price et al., 2011; Vieli and Nick, 2011; Zwally et al., 2011) and this is recognized as a significant limitation in our capacity to accurately predict near-future sea level rise (IPCC, 2007). We therefore highlight numerical modelling as an important area for future development and emphasize the need to combine results with remotely sensed and observational data, in order to improve our understanding of recent changes in Arctic marine-terminating outlet glacier dynamics.

## VII Conclusions

Arctic ice masses have rapidly lost mass since the mid-1990s due to a combination of negative SMB and accelerated discharge from marine-terminating glaciers (van den Broeke et al., 2009). Studies conducted during the past 20

years have fundamentally altered our understanding of ice mass response to climatic/oceanic forcing and have demonstrated that changes in marine-terminating glacier dynamics can result in dramatic mass losses at annual timescales (e.g. Howat et al., 2008b; Rignot and Kanagaratnam, 2006; Stearns and Hamilton, 2007). In this paper, we identify and review three primary climatic/oceanic drivers of marine-terminating Arctic outlet glacier behaviour: air temperatures, ocean temperatures and sea ice. Although discussed separately, these factors are interconnected and we highlight a number of potentially important linkages which may significantly influence glacier dynamics. We suggest that meltwater-enhanced basal sliding may contribute to marine-terminating outlet glacier velocities at seasonal timescales (Howat et al., 2010; Nick et al., 2012), but its net effect on interannual behaviour may be limited, potentially due to the capacity of the subglacial hydrological system to evolve in response to meltwater inputs (Price et al., 2008; Sundal et al., 2011). Instead, marine-terminating outlet glaciers may respond to atmospheric warming via a number of alternative mechanisms, including: (1) hydrofracture of crevasses at the terminus/lateral margins; (2) meltwater-enhanced submarine melting, via plume circulation; and (3) sea ice loss due to atmospheric warming. Marine-terminating outlet glaciers are potentially highly sensitive to oceanic warming (Rignot et al., 2010), which may cause retreat through: (1) submarine melting and rapid thinning across floating sections; (2) grounding-line retreat; (3) alteration of the calving front geometry at the grounding line and/or waterline; and (4) sea ice loss due to oceanic warming. We emphasize the need to further investigate controls on Atlantic Water distribution within glacier fjords and feedbacks between fjord circulation, subglacial meltwater and submarine melting. We also underscore the influence of sea ice on seasonal and interannual outlet glacier dynamics, via its influence on calving rates

(Amundson et al., 2010; Joughin et al., 2008c), and suggest that sea ice forcing may become increasingly important during the 21st century if current negative trends continue.

We suggest that the respective role of each climatic/oceanic factor varies across the Arctic and that outlet glacier response to forcing within one region cannot be assumed to apply elsewhere. Moreover, glacier-specific factors may substantially modulate the response of individual glaciers to climatic/oceanic forcing and we highlight this as priority area for future research. Numerical modelling results have improved our understanding of marine-terminating outlet glacier behaviour, but remain a key area for future development. Notwithstanding recent advances, substantial uncertainties remain over the respective roles of the various climatic/oceanic and glacier-specific forcing factors, and we highlight the potential danger of extrapolating mass loss rates from a small number of study glaciers. Consequently, the response of marine-terminating Arctic outlet glaciers to climatic/oceanic forcing remains a key area for future research and is crucial for accurate prediction of near-future sea level rise and Arctic ice mass response to climate warming.

### Acknowledgements

We thank T. Moon for the provision of Greenland frontal position data and G. Moholdt for helpful comments on Svalbard mass balance data. We thank E. Hanna and N.J. Cox for their assistance with the air temperature data. Two anonymous reviewers are thanked for their constructive comments on the manuscript.

### Funding

This work was supported by a Durham Doctoral Studentship granted to J.R. Carr.

### References

- Abdalati W and Steffen K (2001) Greenland ice sheet melt extent: 1979–1999. *Journal of Geophysical Research* 106: 33983–33988.



- Abdalati W, Krabill W, Frederick E, et al. (2001) Outlet glacier and margin elevation changes: Near-coastal thinning of the Greenland ice sheet. *Journal of Geophysical Research* 106: 33729–33741.
- Abdalati W, Krabill W, Frederick E, et al. (2004) Elevation changes of ice caps in the Canadian Arctic Archipelago. *Journal of Geophysical Research* 109: F04007.
- Ahn Y and Box JE (2010) Glacier velocities from time-lapse photos: Technique development and first results from the Extreme Ice Survey (EIS) in Greenland. *Journal of Glaciology* 198: 723–734.
- Alley RB (1991) Sedimentary processes may cause fluctuations of tidewater glaciers. *Annals of Glaciology* 15: 119–124.
- Alley RB, Dupont TK, Parisek BR, et al. (2005) Access of surface meltwater to beds of sub-freezing glaciers: Preliminary insights. *Annals of Glaciology* 40: 8–14.
- Amundson JM, Fahnestock M, Truffer M, et al. (2010) Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland. *Journal of Geophysical Research* 115: F01005.
- Andersen CS, Larsen TB, Nettles M, et al. (2010a) Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics. *Journal of Geophysical Research* 115: F04041.
- Andersen ML, Larsen TB, Nettles M, et al. (2010b) Spatial and temporal melt variability at Helheim Glacier, East Greenland, and its effect on ice dynamics. *Journal of Geophysical Research* 115: F04041.
- Andersen CS, Straneo F, Hvid M, et al. (2012) Rapid response of Helheim Glacier in Greenland to climate variability over the past century. *Nature Geoscience* 5: 37–41.
- Arctic Climate Impact Assessment (ACIA) (2004) *Impacts of a Warming Arctic: Arctic Climate Impact Assessment*. Cambridge: Cambridge University Press.
- Arendt A, Echelmeyer K, Harrison W, et al. (2006) Updated estimates of glacier volume changes in the western Chugach Mountains, Alaska, and a comparison of regional extrapolation methods. *Journal of Geophysical Research* 111: F03019.
- Arendt A, Luthcke SB, Larsen CF, et al. (2008) Validation of high-resolution GRACE mascon estimates of glacier mass changes in the St Elias Mountains, Alaska, USA, using aircraft laser altimetry. *Journal of Glaciology* 54: 778–787.
- Bamber JL, Alley RB, and Joughin I (2007) Rapid response of modern day ice sheets to external forcing. *Earth and Planetary Science Letters* 257: 1–13.
- Bamber J, Krabill W, Raper V, et al. (2004) Anomalous recent growth of part of a large Arctic ice cap: Austfonna, Svalbard. *Geophysical Research Letters* 31: L12402.
- Bartholomew I, Nienow P, Mair D, et al. (2010) Seasonal evolution of subglacial drainage and acceleration in a Greenland outlet glacier. *Nature Geoscience* 3: 408–411.
- Bartholomew I, Sole A, Mair D, et al. (2011) Supraglacial forcing of subglacial drainage in the ablation zone of the Greenland ice sheet. *Geophysical Research Letters* 38: L08502.
- Bassford RP, Siegert MJ, Dowdeswell J, et al. (2006) Quantifying the mass balance of ice caps on Severnaya Zemlya, Russian High Arctic. I: Climate and mass balance of the Vavilov Ice Cap. *Arctic, Antarctic, and Alpine Research* 38: 1–12.
- Benn DI and Evans DJA (2010) *Glaciers and Glaciation*. London: Hodder Education.
- Benn DI, Warren CR, and Mottram RH. (2007) Calving processes and the dynamics of calving glaciers. *Earth Science Reviews* 82: 143–179.
- Bergmann I, Ramillien G, and Frappart F (2012) Climate-driven interannual ice mass evolution in Greenland. *Global and Planetary Change* 82–83: 1–11.
- Bevan SL, Luckman AJ, and Murray T (2012a) Glacier dynamics over the last quarter of a century at Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers. *The Cryosphere* 6: 923–937.
- Bevan SL, Luckman A, Murray T, et al. (2007) Positive mass balance during the late 20th century on Austfonna, Svalbard, revealed using satellite radar interferometry. *Annals of Glaciology* 46: 117–122.
- Bevan SL, Murray T, Luckman A, et al. (2012b) Stable dynamics in a Greenland tidewater glacier over 26 years despite reported thinning. *Annals of Glaciology* 53: 241–248.
- Bhattacharya I, Jezek KC, Wang L, et al. (2009) Surface melt area variability of the Greenland ice sheet: 1979–2008. *Geophysical Research Letters* 36: L20502.
- Bingham RG, Hubbard A, Nienow P, et al. (2008) An investigation into the mechanisms controlling seasonal speedup events at a High Arctic glacier. *Journal of Geophysical Research* 113: F02006.
- Bingham RG, Nienow P, and Sharp MJ (2003) Intra-annual and intra-seasonal flow dynamics of a High Arctic polythermal valley glacier. *Annals of Glaciology* 37: 181–188.
- Bingham RG, Nienow P, Sharp MJ, et al. (2005) Subglacial drainage processes at a High Arctic polythermal valley glacier. *Journal of Glaciology* 51: 15–24.

- Bingham RG, Nienow P, Sharp MJ, et al. (2006) Hydrology and dynamics of a polythermal (mostly cold) High Arctic glacier. *Earth Surface Processes and Landforms* 31: 1463–1479.
- Bjørk AA, Kjær KH, Korsgaard NJ, et al. (2012) An aerial view of 80 years of climate-related glacier fluctuations in southeast Greenland. *Nature Geoscience* 5: 427–432.
- Błaszczyc M, Jania JA, and Hagen JM (2009) Tidewater glaciers of Svalbard: Recent changes and estimates of calving fluxes. *Polish Polar Research* 30: 85–142.
- Bond G, Heinrich H, Broecker W, et al. (1992) Evidence for massive discharges of icebergs into the North Atlantic ocean during the last glacial period. *Nature* 360: 245–249.
- Boon S and Sharp MJ (2003) The role of hydrologically-driven ice fracture in drainage system evolution on an Arctic glacier. *Geophysical Research Letters* 30: 1916.
- Box JE and Ski K (2007) Remote sounding of Greenland supraglacial melt lakes: Implications to sub-glacial hydraulics. *Journal of Glaciology* 181: 257–265.
- Box JE, Bromwich DH, Veenhuis BA, et al. (2006) Greenland Ice Sheet surface mass balance variability (1988–2004) from calibrated Polar MM5 output. *Journal of Climate* 19: 2783–2800.
- Box JE, Yang L, Bromwich DH, et al. (2009) Greenland Ice Sheet surface air temperature variability: 1840–2007. *Journal of Climate* 22: 4029–4049.
- Briner JP, Bini AC, and Anderson RS (2009) Rapid early Holocene retreat of a Laurentide outlet glacier through an Arctic fjord. *Nature Geoscience* 2: 496–499.
- Burgess DO and Sharp MJ (2004) Recent changes in areal extent of the Devon Ice Cap, Nunavut, Canada. *Arctic, Antarctic, and Alpine Research* 36: 261–271.
- Burgess DO and Sharp MJ (2008) Recent changes in thickness of the Devon Island ice cap, Canada. *Journal of Geophysical Research* 113: B07204.
- Cappelen J (ed.) (2011) Technical Report 11-05: DMI monthly climate data collection 1768–2010, Denmark, The Faroe Islands and Greenland. Copenhagen: Danish Meteorological Institute.
- Catania GA and Neumann TA (2010) Persistent englacial drainage features in the Greenland Ice Sheet. *Geophysical Research Letters* 37: L02501.
- Christoffersen P, Mugford R, Heywood KJ, et al. (2011) Warming of waters in an East Greenland fjord prior to glacier retreat: Mechanisms and connection to large-scale atmospheric conditions. *The Cryosphere Discussions* 5: 1335–1364.
- Chylek P, Dubey MK and Lesins G (2006) Greenland warming of 1920–1930 and 1995–2005. *Geophysical Research Letters* 33: L11707.
- Copland L, Sharp MJ, and Nienow P (2003) Links between short-term velocity variations and the subglacial hydrology of a predominantly cold polythermal glacier. *Journal of Glaciology* 49: 337–348.
- Csatho B, Schenk T, van der Veen CJ, et al. (2008) Intermittent thinning of Jakobshavn Isbræ, West Greenland, since the Little Ice Age. *Journal of Glaciology* 54: 131–144.
- Das SB, Joughin I, Behn MD, et al. (2008) Fracture propagation to the base of the Greenland Ice Sheet during supraglacial lake drainage. *Science* 320: 778–781.
- Dowdeswell J, Benham TJ, Gorman MR, et al. (2004) Form and flow of the Devon Island Ice Cap, Canadian Arctic. *Journal of Geophysical Research* 109: F02002.
- Dowdeswell J, Benham TJ, Strozzi T, et al. (2008) Iceberg calving flux and mass balance of the Austfonna ice cap on Nordaustlandet, Svalbard. *Journal of Geophysical Research* 113: F03022.
- Dowdeswell J, Hagen JM, Björnsson H, et al. (1997) The mass balance of circum-Arctic glaciers and recent climate change. *Quaternary Research* 48: 1–14.
- Dunse T, Schuler TV, Hagen JO, et al. (2012) Seasonal speed-up of two outlet glaciers of Austfonna, Svalbard, inferred from continuous GPS measurements. *The Cryosphere* 6: 453–466.
- Dyrgerov MB and McCabe GJ (2006) Associations between accelerated glacier mass wastage and increased summer temperature in coastal regions. *Arctic, Antarctic, and Alpine Research* 38: 190–197.
- Ettema J, van den Broeke M, van Meijgaard E, et al. (2009) Higher surface mass balance of the Greenland Ice Sheet revealed by high-resolution climate modeling. *Geophysical Research Letters* 36: L12501.
- Flato GM (2004) Sea-ice and its response to CO<sub>2</sub> forcing as simulated by global climate models. *Climate Dynamics* 23: 229–241.
- Fountain AG and Walder JS (1998) Water flow through temperate glaciers. *Reviews of Geophysics* 36: 299–328.
- Gardner S, Moholdt G, Wouters B, et al. (2011) Sharply increased mass loss from glaciers and ice caps in the Canadian Arctic Archipelago. *Nature* 473: 357–360.

- Gerdes R, Larcher MJ, Kauker F, et al. (2003) Causes and development of repeated Arctic Ocean warming events. *Geophysical Research Letters* 30: OCE2.1–2.4.
- Greve R (2000) On the response of the Greenland Ice Sheet to greenhouse climate change. *Climatic Change* 46: 289–303.
- Gudmundsson GH, Krug J, Durand G, et al. (2012) The stability of grounding lines on retrograde slopes. *The Cryosphere Discussions* 6: 2597–2619.
- Hagen JM, Dunse T, Eiken T, et al. (2009) GLACIODYN – the dynamic response of Arctic glaciers to global warming. American Geophysical Union, Fall Meeting. Abstract C53B-06.
- Hall BL, Baroni C, Denton GH, et al. (2008) Relative sea-level change, Kjove Land, Scoresby Sund, East Greenland: Implications for seasonality in late-glacial time. *Quaternary Science Reviews* 27: 2283–2291.
- Hanna E, Cappelen J, Fettweis X, et al. (2009) Hydrologic response of the Greenland Ice Sheet: The role of oceanographic warming. *Hydrological Processes* 23: 7–30.
- Hanna E, Huybrechts P, Steffen K, et al. (2008) Increased runoff from melt from the Greenland Ice Sheet: A response to global warming. *Journal of Climate* 21: 331–341.
- Higgins AK (1989) North Greenland ice islands. *Polar Record* 25: 209–212.
- Higgins AK (1990) North Greenland glacier velocities and calf ice production. *Polarforschung* 60: 1–23.
- Hodgkins R (1997) Glacier hydrology in Svalbard, Norwegian High Arctic. *Quaternary Science Reviews* 16: 957–973.
- Holland DM, Thomas RH, de Young B, et al. (2008) Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters. *Nature Geoscience* 1: 1–6.
- Holliday NP, Hughes SL, Bacon S, et al. (2008) Reversal of the 1960s to 1990s freshening trend in the northeast North Atlantic and Nordic Seas. *Geophysical Research Letters* 35: L03614.
- Howat IM and Eddy A (2011) Multi-decadal retreat of Greenland's marine-terminating glaciers. *Journal of Glaciology* 57: 389–396.
- Howat IM, Ahn Y, Joughin I, et al. (2011) Mass balance of Greenland's three largest outlet glaciers, 2000–2010. *Geophysical Research Letters* 38: L12501.
- Howat IM, Box JE, Ahn Y, et al. (2010) Seasonal variability in the dynamics of marine-terminating outlet glaciers in Greenland. *Journal of Glaciology* 56: 601–613.
- Howat IM, Joughin I, Fahnestock M, et al. (2008a) Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–2006; Ice dynamics and coupling to climate. *Journal of Glaciology* 54: 1–14.
- Howat IM, Joughin I, and Scambos TA (2007) Rapid changes in ice discharge from Greenland outlet glaciers. *Science* 315: 1559–1561.
- Howat IM, Joughin I, Tulaczyk S, et al. (2005) Rapid retreat and acceleration of Helheim Glacier, east Greenland. *Geophysical Research Letters* 32: L22502.
- Howat IM, Smith BE, Joughin I, et al. (2008b) Rates of southeast Greenland ice volume loss from combined ICESat and ASTER observations. *Geophysical Research Letters* 35: L17505.
- Hughes NE, Wilkinson JP, and Wadhams P (2011) Multi-satellite sensor analysis of fast-ice development in the Norske Øer Ice Barrier, northeast Greenland. *Annals of Glaciology* 52: 151–160.
- Hughes T (1986) The Jakobshavns effect. *Geophysical Research Letters* 13: 46–48.
- Hurrell JW, Kushnir Y, Visbeck MM, et al. (2003) An overview of the North Atlantic Oscillation. In: Hurrell JW, Kushnir Y, and Ottersen GG, et al. (eds) *The North Atlantic Oscillation: Climate Significance and Environmental Impact*. Geophysical Monograph Series 134. Washington, DC: American Geophysical Union, 1–35.
- Huybrechts P and de Wolde J (1999) The dynamic response of the Greenland and Antarctic ice sheets to multiple-century warming. *Journal of Climate* 21: 2169–2188.
- Huybrechts P, Letreuilly A, and Reeh N (1991) The Greenland ice sheet and greenhouse warming. *Palaeogeography, Palaeoclimatology, Palaeoecology* 89: 399–412.
- Iken A and Bindshadler RA (1986) Combined measurements of subglacial water pressure and surface velocity of Findelengletscher, Switzerland: Conclusions about drainage system and sliding mechanism. *Journal of Glaciology* 32: 101–119.
- Intergovernmental Panel on Climate Change (IPCC) (2001) *The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC) (2007) *The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of*

- the Intergovernmental Panel on Climate Change. Cambridge: Cambridge University Press.
- Jacob T, Wahr J, Pfeffer WT, et al. (2012) Recent contributions of glaciers and ice caps to sea level rise. *Nature* 482: 514–518.
- Jamieson SSR, Vieli A, Livingstone SJ, et al. (2012) Ice stream stability on a reverse bed slope. *Nature Geoscience* 5: 799–802.
- Jenkins A (2011) Convection-driven melting near the grounding lines of ice shelves and tidewater glaciers. *Journal of Physical Oceanography* 41: 2279–2294.
- Johannessen OM, Khvorostovsky K, Miles MW, et al. (2005) Recent ice-sheet growth in the interior of Greenland. *Science* 310: 1013–1016.
- Johnson HL, Münchow A, Falkner KK, et al. (2011) Ocean circulation and properties in Petermann Fjord, Greenland. *Journal of Geophysical Research* 116: C01003.
- Joughin I and Alley RB (2011) Stability of the West Antarctic ice sheet in a warming world. *Nature Geoscience* 4: 506–513.
- Joughin I, Abdalati W, and Fahnestock M (2004) Large fluctuations in speed on Greenland's Jakobshavn Isbrae glacier. *Nature* 432: 608–610.
- Joughin I, Das SB, King MA, et al. (2008a) Seasonal speedup along the western flank of the Greenland Ice Sheet. *Science* 320: 781–783.
- Joughin I, Howat IM, Alley RB, et al. (2008b) Ice-front variation and tidewater behaviour on Helheim and Kangerdlugssuaq Glaciers, Greenland. *Journal of Geophysical Research* 113: F01004.
- Joughin I, Howat IM, Fahnestock M, et al. (2008c) Continued evolution of Jakobshavn Isbrae following its rapid speedup. *Journal of Geophysical Research* 113: F04006.
- Joughin I, Smith B, Howat IM, et al. (2010) Greenland flow variability from ice-sheet-wide velocity mapping. *Journal of Glaciology* 56: 415–430.
- Joughin I, Smith BE, Howat I, et al. (2012) Seasonal to decadal scale variations in the surface velocity of Jakobshavn Isbrae, Greenland: Observation and model-based analysis. *Journal of Geophysical Research* 117: F02030.
- Kamb B (1987) Glacier surge mechanism based on linked cavity configuration of the basal water conduit system. *Journal of Geophysical Research* 92: 9083–9100.
- Khan SA, Wahr J, Bevis M, et al. (2010) Spread of ice mass loss into northwest Greenland observed by GRACE and GPS. *Geophysical Research Letters* 37: L06501.
- Kjær KH, Khan SA, Korsgaard NJ, et al. (2012) Aerial photographs reveal late-20th-century dynamic ice loss in northwestern Greenland. *Science* 337: 596–573.
- Koerner R (2005) Mass balance of glaciers in the Queen Elizabeth Islands, Nunavut, Canada. *Annals of Glaciology* 42: 417–423.
- Kotlyakov VM, Glazovskii AF, and Frolov IE (2010) Glaciation in the Arctic. *Herald of the Russian Academy of Sciences* 80: 155–164.
- Krabill W, Hanna E, Huybrechts P, et al. (2004) Greenland Ice Sheet: Increased coastal thinning. *Geophysical Research Letters* 31: L24402.
- Krawczynski MJ, Behn MD, Das SB, et al. (2009) Constraints on the lake volume required for hydro-fracture through ice sheets. *Geophysical Research Letters* 36: L10501.
- Kwok R and Rothcock DA (2009) Decline in Arctic sea ice thickness from submarine and ICESat records: 1958–2008. *Geophysical Research Letters* 36: L15501.
- Luckman A and Murray T (2005) Seasonal variation in velocity before retreat of Jakobshavn Isbrae, Greenland. *Geophysical Research Letters* 32: L08501.
- Luckman A, Murray T, de Lange R, et al. (2006) Rapid and synchronous ice-dynamic changes in East Greenland. *Geophysical Research Letters* 33: L03503.
- Luthcke SB, Zwally HJ, Abdalati W, et al. (2006) Recent Greenland ice mass loss by drainage system from satellite gravity observations. *Science* 314: 1286–1289.
- McFadden EM, Howat IM, Joughin I, et al. (2011) Changes in the dynamics of marine terminating outlet glaciers in west Greenland (2000–2009). *Journal of Geophysical Research* 116: F02022.
- McMillan M, Nienow P, Shepherd A, et al. (2007) Seasonal evolution of supra-glacial lakes on the Greenland ice sheet. *Earth Planetary Science Letters* 262: 484–492.
- Mair D, Burgess DO, Sharp MJ, et al. (2009) Mass balance of the Prince of Wales Icefield, Ellesmere Island, Nunavut, Canada. *Journal of Geophysical Research* 114: F02011.
- Mayer C, Reeh N, Jung-Rothenhäusler F, et al. (2000) The subglacial cavity and implied dynamics under Niothalvfjærdsfjorden Glacier, NE-Greenland. *Geophysical Research Letters* 27: 2289–2292.
- Meier MF and Post A (1987) Fast tidewater glaciers. *Journal of Geophysical Research* 92: 9051–9058.

- Meier MF, Dyurgerov MB, Rick UK, et al. (2007) Glaciers dominate eustatic sea-level rise in the 21st century. *Science* 317: 1064–1067.
- Mémin A, Rogister Y, Hinderer J, et al. (2011) Secular gravity variation at Svalbard (Norway) from ground observations and GRACE satellite data. *Geophysical Journal International* 184: 1119–1130.
- Moholdt G, Hagen JM, Eiken T, et al. (2010a) Geometric changes and mass balance of the Austfonna ice cap, Svalbard. *The Cryosphere* 4: 21–34.
- Moholdt G, Nuth C, Hagen JO, et al. (2010b) Recent elevation changes of Svalbard glaciers derived from ICESat laser altimetry. *Remote Sensing of Environment* 114: 2756–2767.
- Moholdt G, Wouters B, and Gardner AS (2012) Recent mass changes of glaciers in the Russian High Arctic. *Geophysical Research Letters* 39: L10502.
- Moon T and Joughin I (2008) Changes in ice-front position on Greenland's outlet glaciers from 1992 to 2007. *Journal of Geophysical Research* 113: F02022.
- Moon T, Joughin I, Smith BE, et al. (2012) 21st-century evolution of Greenland outlet glacier velocities. *Science* 336: 576–578.
- Mortensen J, Lennert K, Bendtsen J, et al. (2011) Heat sources for glacial melt in a sub-Arctic fjord (Godthåbsfjord) in contact with the Greenland Ice Sheet. *Journal of Geophysical Research* 116: C01013.
- Mote TL (2007) Greenland surface melt trends 1973–2007: Evidence of a large increase in 2007. *Geophysical Research Letters* 34: L22507.
- Motyka RJ, Fahnestock M, and Truffer M (2010) Volume change of Jakobshavn Isbræ, West Greenland: 1985–1997–2007. *Journal of Glaciology* 56: 635–645.
- Motyka RJ, Hunter L, Echelmeyer K, et al. (2003) Submarine melting at the terminus of a temperate tidewater glacier, LeConte Glacier, Alaska, USA. *Annals of Glaciology* 36: 57–65.
- Motyka RJ, Truffer M, Fahnestock M, et al. (2011) Submarine melting of the 1985 Jakobshavn Isbræ floating tongue and the triggering of the current retreat. *Journal of Geophysical Research* 116: F01007.
- Murray T, Scharrer K, James TD, et al. (2010) Ocean regulation hypothesis for glacier dynamics in southeast Greenland and implications for ice sheet mass changes. *Journal of Geophysical Research* 115: F03026.
- Nick FM, Luckman A, Vieli A, et al. (2012) The response of Petermann Glacier, Greenland, to large calving events, and its future stability in the context of atmospheric and oceanic warming. *Journal of Glaciology* 58: 229–239.
- Nick FM, van der Veen CJ, Vieli A, et al. (2010) A physically based calving model applied to marine outlet glaciers and implications for the glacier dynamics. *Journal of Geophysical Research* 115: 781–794.
- Nick FM, Vieli A, Howat IM, et al. (2009) Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus. *Nature Geoscience* 2: 110–114.
- Nienow P, Sharp MJ, and Willis IC (1998) Seasonal changes in the morphology of the subglacial drainage system, Haut Glacier d'Arolla, Switzerland. *Earth Surface Processes and Landforms* 23: 825–843.
- Nilsen F, Cottier F, Skogseth R, et al. (2008) Fjord-shelf exchanges controlled by ice and brine production: The interannual variation of Atlantic Water in Isfjorden, Svalbard. *Continental Shelf Research* 28: 1838–1853.
- Nuth C, Moholdt G, Kohler J, et al. (2010) Svalbard glacier elevation changes and contribution to sea level rise. *Journal of Geophysical Research* 115: F01008.
- Nuttall A-M and Hodgkins R (2005) Temporal variations in flow velocity at Finsterwalderbreen, a Svalbard surge-type glacier. *Annals of Glaciology* 42: 71–76.
- Palmer S, Shepherd A, Nienow P, et al. (2011) Seasonal speedup of the Greenland Ice Sheet linked to routing of surface water. *Earth and Planetary Science Letters* 302: 423–428.
- Parisek BR and Alley RB (2004) Implications of increased Greenland surface melt under global-warming scenarios: Ice-sheet simulations. *Quaternary Science Reviews* 23: 1013–1027.
- Payne AJ, Vieli A, Shepherd AP, et al. (2004) Recent dramatic thinning of largest West Antarctic ice stream triggered by oceans. *Geophysical Research Letters* 31: L23401.
- Pfeffer WT (2003) Tidewater glaciers move at their own pace. *Nature* 426: 602.
- Pfeffer WT (2007) A simple mechanism for irreversible tidewater glacier retreat. *Journal of Geophysical Research* 112: F03S25.
- Price S, Conway H, Waddington ED, et al. (2008) Model investigations of inland migration of fast-flowing outlet glaciers and ice streams. *Journal of Glaciology* 54: 49–60.
- Price S, Payne AJ, Howat IM, et al. (2011) Committed sea-level rise for the next century from Greenland ice sheet



- dynamics during the past decade. *Proceedings of the National Academy of Sciences* 108: 8978–8983.
- Pritchard HD, Arthern RJ, Vaughan DG, et al. (2009) Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets. *Nature* 461: 971–975.
- Radić V and Hock R (2011) Regionally differentiated contribution of mountain glaciers and ice caps to future sea-level rise. *Nature Geoscience* 4: 91–94.
- Raper V, Bamber J, and Krabill W (2005) Interpretation of the anomalous growth of Austfonna, Svalbard, a large Arctic ice cap. *Annals of Glaciology* 42: 373–379.
- Reeh N, Thomsen HH, Higgins AK, et al. (2001) Sea ice and the stability of north and northeast Greenland floating glaciers. *Annals of Glaciology* 33: 474–480.
- Rennermalm AK, Smith LC, Stroeve JC, et al. (2009) Does sea ice influence Greenland ice sheet surface-melt? *Environmental Research Letters* 4: 024011.
- Rignot E and Kanagaratnam P (2006) Changes in the velocity structure of the Greenland Ice Sheet. *Science* 311: 986–990.
- Rignot E and Steffen K (2008) Channelized bottom melting and stability of floating ice shelves. *Geophysical Research Letters* 35: L02503.
- Rignot E, Box JE, Burgess E, et al. (2008) Mass balance of the Greenland ice sheet from 1958 to 2007. *Geophysical Research Letters* 35: L20502.
- Rignot E, Fenty I, Menemenlis D, et al. (2012) Spreading of warm ocean waters around Greenland as a possible cause for glacier acceleration. *Annals of Glaciology* 53: 257–266.
- Rignot E, Koppes M, and Velicogna I (2010) Rapid submarine melting of the calving faces of West Greenland glaciers. *Nature Geoscience* 3: 187–191.
- Rignot E, Velicogna I, van den Broeke M, et al. (2011) Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise. *Geophysical Research Letters* 38: L05503.
- Rippin DM, Willis IC, Arnold NS, et al. (2005) Spatial and temporal variations in surface velocity and basal drag across the tongue of the polythermal glacier midre Lovénbreen, Svalbard. *Journal of Glaciology* 51: 588–600.
- Rodrigues J (2008) The rapid decline of the sea ice in the Russian Arctic. *Cold Regions Science and Technology* 54: 124–142.
- Rodrigues J (2009) The increase in the length of the ice-free season in the Arctic. *Cold Regions Science and Technology* 59: 78–101.
- Schoof C (2010) Ice-sheet acceleration driven by melt supply variability. *Nature* 468: 803–806.
- Seale A, Christoffersen P, Mugford R, et al. (2011) Ocean forcing of the Greenland Ice Sheet: Calving fronts and patterns of retreat identified by automatic satellite monitoring of eastern outlet glaciers. *Journal of Geophysical Research* 116: F03013.
- Serreze MC, Barrett AP, Stroeve JC, et al. (2009) The emergence of surface-based Arctic amplification. *The Cryosphere* 3: 11–19.
- Sharov AI (2005) Studying changes of ice coasts in the European Arctic. *Geo-Marine letters* 25: 153–166.
- Sharov AI (2010) *Satellite Monitoring and Regional Analysis of Glacier Dynamics in the Barents-Kara Region*. Graz: Joanneum Research Forschungsgesellschaft mbH, Institute of Digital Image Processing.
- Sharov AI, Schöner W, and Pail R (2009) Spatial features of glacier changes in the Barents-Kara Sector. *EGU General Assembly* 11: EGU2009–3046.
- Shepherd A, Du Z, Benham TJ, et al. (2007) Mass balance of Devon Ice Cap, Canadian Arctic. *Annals of Glaciology* 46: 249–254.
- Shepherd A, Hubbard A, Nienow P, et al. (2009) Greenland ice sheet motion coupled with daily melting in late summer. *Geophysical Research Letters* 36: L01501.
- Sneed WA and Hamilton GS (2007) Evolution of melt pond volume on the surface of the Greenland Ice Sheet. *Geophysical Research Letters* 34: L03501.
- Sohn HG, Jezek KC, and van der Veen CJ (1998) Jakobshavn Glacier, West Greenland: 30 years of spaceborne observations. *Geophysical Research Letters* 25: 2699–2702.
- Sole AJ, Mair DW, Nienow PW, et al. (2011) Seasonal speedup of a Greenland marine-terminating outlet glacier forced by surface melt-induced changes in subglacial hydrology. *Journal of Geophysical Research* 116: F03014.
- Sole A, Payne T, Bamber J, et al. (2008) Testing hypotheses of the cause of peripheral thinning of the Greenland Ice Sheet: Is land-terminating ice thinning at anomalously high rates? *The Cryosphere Discussions* 2: 673–710.
- Stearns A and Hamilton GS (2007) Rapid volume loss from two East Greenland outlet glaciers quantified using repeat stereo satellite imagery. *Geophysical Research Letters* 34: L05503.
- Stern H and Heide-Jørgensen MP (2003) Trends and variability of sea ice in Baffin Bay and Davis Strait, 1953–2001. *Polar Research* 22: 11–18.

- Straneo F, Curry RG, Sutherland DA, et al. (2011) Impact of fjord dynamics and glacial runoff on the circulation near Helheim Glacier. *Nature Geoscience* Advanced online publication (10th April 2011).
- Straneo F, Hamilton GS, Sutherland DA, et al. (2010) Rapid circulation of warm subtropical waters in a major glacial fjord in East Greenland. *Nature Geoscience* 3: 182–186.
- Sundal AV, Shepherd A, Nienow P, et al. (2011) Melt-induced speed-up of Greenland ice sheet offset by efficient subglacial drainage. *Nature* 469: 521–524.
- Thomas R (1979) The dynamics of marine ice sheets. *Journal of Glaciology* 24: 167–177.
- Thomas R, Frederick E, Li J, et al. (2011) Accelerating ice loss from the fastest Greenland and Antarctic glaciers. *Geophysical Research Letters* 38: L10502.
- Thomas RH (2004) Force-perturbation analysis of recent thinning and acceleration of Jakobshavn Isbrae, Greenland. *Journal of Glaciology* 50: 57–66.
- Thomas RH, Abdalati W, Frederick E, et al. (2003) Investigation of surface melting and dynamic thinning on Jakobshavn Isbrae, Greenland. *Journal of Glaciology* 49: 231–239.
- Thomas RH, Frederick E, Krabill W, et al. (2006) Progressive increase in ice loss from Greenland. *Geophysical Research Letters* 33: L10503.
- Thomas RH, Frederick E, Krabill W, et al. (2008) A comparison of Greenland ice-sheet volume changes derived from altimetry measurements. *Journal of Glaciology* 54: 203–212.
- Thomas RH, Frederick E, Krabill W, et al. (2009) Recent changes on Greenland outlet glaciers. *Journal of Glaciology* 55: 147–162.
- van den Broeke M, Bamber J, Ettema J, et al. (2009) Partitioning recent Greenland mass loss. *Science* 326: 984–986.
- van der Veen CJ (1998) Fracture mechanics approach to penetration of surface crevasses on glaciers. *Cold Regions Science and Technology* 27: 31–47.
- van der Veen CJ (2007) Fracture propagation as means of rapidly transferring surface meltwater to the base of glaciers. *Geophysical Research Letters* 34: L01501.
- van der Veen CJ, Plummer JC, and Stearns LA (2011) Controls on the recent speed-up of Jakobshavn Isbræ, West Greenland. *Journal of Glaciology* 57: 770–782.
- van de Wal RSW and Oerlemans J (1997) Modelling the short-term response of the Greenland ice-sheet to global warming. *Climate Dynamics* 13: 733–744.
- van de Wal RSW, Boot W, van den Broeke MR, et al. (2008) Large and rapid melt-induced velocity changes in the ablation zone of the Greenland Ice Sheet. *Science* 321: 111–113.
- Velicogna I (2009) Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE. *Geophysical Research Letters* 36: L19503.
- Velicogna I and Wahr J (2006) Acceleration of Greenland ice mass loss in spring 2004. *Nature* 443: 329–331.
- Vieli A and Nick FM (2011) Understanding and modelling rapid dynamic changes of tidewater outlet glaciers: Issues and implications. *Surveys in Geophysics* 32: 437–485.
- Vieli A, Funk M, and Blatter H (2001) Flow dynamics of tidewater glaciers: A numerical modelling approach. *Journal of Glaciology* 47: 595–606.
- Vieli A, Jania JA, Blatter H, et al. (2004) Short-term velocity variations on Hansbreen, a tidewater glacier in Spitsbergen. *Journal of Glaciology* 50: 389–398.
- Vieli A, Jania JA, and Lezek K (2002) The retreat of a tidewater glacier: Observations and model calculations on Hansbreen, Spitsbergen. *Journal of Glaciology* 48: 592–600.
- Vieli A, Payne AJ, Shepherd A, et al. (2007) Causes of pre-collapse changes of the Larsen B ice shelf: Numerical modelling and assimilation of satellite observations. *Earth and Planetary Science Letters* 259: 297–306.
- Weertman J (1974) Stability of the junction of an ice sheet and an ice shelf. *Journal of Glaciology* 13: 3–11.
- Weidick A (1975) *A Review of Quaternary Investigations in Greenland*. Institute of Polar Studies Report 55. Columbus, OH: Ohio State University Research Foundation.
- Williamson S, Sharp MJ, Dowdeswell J, et al. (2008) Iceberg calving rates from northern Ellesmere Island ice caps, Canadian Arctic, 1999–2003. *Journal of Glaciology* 54: 391–400.
- Willis IC (1995) Intra-annual variations in glacier motion: A review. *Progress in Physical Geography* 19: 61–106.
- Winsborrow MCM, Andreassen K, Corner GD, et al. (2010) Deglaciation of a marine-based ice sheet: Late Weichselian palaeo-ice dynamics and retreat in the southern Barents Sea reconstructed from onshore and offshore glacial geomorphology. *Quaternary Science Reviews* 29: 424–442.

- Wouters B, Chambers D, and Schrama EJO (2008) GRACE observes small-scale mass loss in Greenland. *Geophysical Research Letters* 35: L20501.
- Xu Y, Rignot E, Menemenlis D, et al. (2012) Numerical experiments on subaqueous melting of Greenland tide-water glaciers in response to ocean warming and enhanced subglacial discharge. *Annals of Glaciology* 53: 229–234.
- Yin J, Overpeck TJ, Griffies SM, et al. (2012) Different magnitudes of projected subsurface ocean warming around Greenland and Antarctica. *Nature Geoscience* 4: 524–528.
- Zwally HJ, Abdalati W, Herring T, et al. (2002) Surface melt-induced acceleration of Greenland ice-sheet flow. *Science* 297: 218–222.
- Zwally HJ, Giovinetto MB, Jun L, et al. (2005) Mass changes of the Greenland and Antarctic ice sheets and shelves and contributions to sea level rise: 1992–2002. *Journal of Glaciology* 51: 509–527.
- Zwally HJ, Li J, Brenner AC, et al. (2011) Greenland ice sheet mass balance: Distribution of increased mass loss with climate warming; 2003–07 versus 1992–2002. *Journal of Glaciology* 57: 88–102.