

Fever evaluation, basic

Authors: Kaur G, Ahmed S, Johansson KA, Coates MM, Watkins D, Økland JM, Haaland ØA.

Date: 2021-07-31

Date updated: 2021-11-20

Description of condition and intervention

Fever evaluation is an important component of routine health care. Presenting with fever is a common symptom reported by individuals for seeking health care. Causative agents for fever could range from infections caused by like bacterial, viral, protozoal, fungal organisms or non-infectious causes. Two guidelines by World Health Organization on Integrated Management of Adolescent and Adult Illness (IMAI) guidelines for health workers at first level facilities (health centers and first-level) (WHO 2009), and also at district hospitals (WHO 2011) provide guidance to health professionals about management of febrile illnesses including fever, especially in resource-limited settings.

In this evidence brief, we present the effect and cost of the following intervention being analysed in FairChoices:DCP Analytical tool:

Pneumonia, oral antibiotics

Diarrhea, oral rehydration therapy (ORT) and zinc

*Malaria, rapid diagnostic tests and antimalarials (principally artemisinin-based combination therapy)**

International guidelines

Organization	Indications/recommendations	Applicability in LIC & Lower MIC settings
WHO 2009	Integrated Management of Adolescent and Adult Illness (IMAI). Guidelines for first-level facility health workers at health centre and district outpatient clinic	Yes

WHO 2011	IMAI district clinician manual: hospital care adolescents and adults: guidelines for the management of illnesses with limited-resources	Yes
----------	---	-----

Intervention attributes

Type of interventions

Table 1: Type of interventions & delivery platform

Intervention	Type	Delivery platform
Pneumonia, oral antibiotics	Curative	Health centre
Diarrhoea, oral rehydration therapy (ORT) and zinc	Curative	Health centre
Malaria, rapid diagnostic tests and antimalarials (principally artemisinin-based combination therapy) *	Curative	Health centre

*This intervention is considered for effects and costs in Malaria cluster

Equity

In addition to considerations like cost-effectiveness and health systems factors, dimensions of equity can be relevant for priority setting. The opportunity for a long and healthy life varies according to the severity of a health condition that individuals might have, so there are inequities in individuals' opportunities for long and healthy lives based on the health conditions they face. Metrics used to estimate the severity of illness at an individual level can be used to help prioritize those with less opportunity for lifetime health. FairChoices: DCP Analytics Tool uses Health adjusted age of death (HAAD), which is a metric that estimates the number of years lived from birth to death, discounting years lived with disability. A high HAAD thus represents a disease less severe in terms of lifetime health loss, while a low HAAD represents a disease that is severe on average, causing early death or a long period of severe disability. It is also possible to estimate the distribution of HAAD across individuals with a health condition. FairChoices shows for each intervention an average HAAD value of the conditions that are affected by respective interventions that have health effects. Additionally, a plot shows HAAD values for around 290 conditions (Johansson KA et al 2020).

(DCP4 ID: INFCTN03-01,02,03)

Cluster: Infection in general

Time dependence

High level of urgency. Treatment outcomes may be affected by some days of delay.

Population in need of interventions

Table 2: Population in need of interventions

Intervention	Treated population		Affected population		Disease state addressed
	Treated age	Treated fraction	Affected age	Affected fraction	
Pneumonia, oral antibiotics	5 to 99 years both genders; incidence based	0.8*	5 to 99	0.8*	For effects: Lower respiratory infections
Diarrhoea, oral rehydration therapy (ORT) and zinc	5 to 99 years both genders; incidence based	0.8*	5 to 99	0.8*	For effects: Diarrheal diseases
Malaria, rapid diagnostic tests and antimalarials (principally artemisinin-based combination therapy) **	**The effects and costs of this intervention are considered in the malaria cluster.				

* Assumed based on expert opinion

Intervention effect and safety

Table 3: Effect and safety of interventions for fever evaluation

Effect of intervention	Certainty of evidence
------------------------	-----------------------

(DCP4 ID: INFCTN03-01,02,03)

Cluster: Infection in general

Mortality (due to condition)		
Pneumonia, oral antibiotics (Amoxicillin efficacy for pneumonia case management in children)	0.7 relative risk reduction) (Johansson KA et al 2020 & Theodoratou E et al 2010)	See appendix
Diarrhea, oral rehydration therapy (ORT)	0.93 (Pecenka CJ et al 2015 & Munos, Walker, Black 2010)	

Model assumptions

Table 2: Summary of model parameters and values used in FairChoices – DCP Analytical Tool

Category	Model parameter	Notes
Interventions	Pneumonia, oral antibiotics Diarrhea, oral rehydration therapy (ORT)	
Cost calculation		
Treated population	See Table 2	Global Burden of disease study 2019
Gender		
Age		
Treated fraction		
Effect calculation		
Affected Population	See Table 2	
Affected gender		
Affected fraction age		
Affected fraction		
Comparison	placebo or other care	

(DCP4 ID: INFCTN03-01,02,03)

Cluster: Infection in general

Mortality Reduction (RRR) Pneumonia, oral antibiotics	0.7	(Johansson KA et al 2020 & Theodoratou E et al 2010)
Diarrhea, oral rehydration therapy	0.93	(Pecenka CJ et al 2015 & Munos, Walker, Black 2010)

Intervention Cost

The cost for fever evaluation (adolescents | adults) and essential management, clinically stable according to the WHO IMAI guidelines, with the referral of unstable individuals, was calculated using the per-incident case for managing typhoid fever, Dengue, and influenza-associated hospitalizations as a proxy with the total cost based on the sum of management of each condition, i.e., 2 USD (india.2004) (Sur D et al 2009), 23.49 USD (India,2012) (Shepard DS et al 2014) and 10.52 USD (Bangladesh,2010) (Bhuiyan MU et al 2014) consecutively. Each cost is divided by three to get the average cost for managing a stable case of fever.

References

WHO. Acute Care. Integrated Management of Adolescent and Adult Illness (IMAI). Guidelines for first-level facility health workers at health centre and district outpatient clinic.2009

World Health Organization. IMAI district clinician manual: hospital care adolescents and adults: guidelines for the management of illnesses with limited-resources. 2011

Johansson KA et al 2020: Johansson KA, Memirie ST, Pecenka C, Jamison DT, Verguet S. Health Gains and Financial Protection from Pneumococcal Vaccination and Pneumonia Treatment in Ethiopia: Results from an Extended Cost-Effectiveness Analysis. PLoS One. 2015 Dec 9;10(12): e0142691. doi: 10.1371/journal.pone.0142691. PMID: 26650078; PMCID: PMC4674114.

Theodoratou E et al 2010: Theodoratou E, Al-Jilaihawi S, Woodward F, Ferguson J, Jhass A, Balliet M, Kolcic I, Sadruddin S, Duke T, Rudan I, Campbell H. The effect of case management on childhood pneumonia mortality in developing countries. Int J Epidemiol. 2010 Apr;39 Suppl 1(Suppl 1):i155-71. doi: 10.1093/ije/dyq032. PMID: 20348118; PMCID: PMC2845871.

(DCP4 ID: INFCTN03-01,02,03)

Cluster: Infection in general

Pecenka CJ et al 2015: Pecenka CJ, Johansson KA, Memirie ST, Jamison DT, Verguet S. Health gains and financial risk protection: an extended cost-effectiveness analysis of treatment and prevention of diarrhoea in Ethiopia. *BMJ Open*. 2015 May 3;5(4):e006402. doi: 10.1136/bmjopen-2014-006402. PMID: 25941175; PMCID: PMC4420944.

Munos, Walker, Black 2010: Munos MK, Walker CL, Black RE. The effect of oral rehydration solution and recommended home fluids on diarrhoea mortality. *Int J Epidemiol*. 2010 Apr;39 Suppl 1(Suppl 1):i75-87. doi: 10.1093/ije/dyq025. PMID: 20348131; PMCID: PMC2845864.

Sur D et al 2009: Sur D, Chatterjee S, Riewpaiboon A, Manna B, Kanungo S, Bhattacharya SK. Treatment cost for typhoid fever at two hospitals in Kolkata, India. *J Health Popul Nutr*. 2009 Dec;27(6):725-32. doi: 10.3329/jhpn.v27i6.4323. PMID: 20099755; PMCID: PMC2928117.

Shepard DS et al 2014: Shepard DS, Halasa YA, Tyagi BK, Adhish SV, Nandan D, Karthiga KS, Chellawamy V, Gaba M, Arora NK, The Inclen Study Group. Economic and disease burden of dengue illness in India. *Am J Trop Med Hyg*. 2014 Dec;91(6):1235-1242. doi: 10.4269/ajtmh.14-0002. Epub 2014 Oct 6. PMID: 25294616; PMCID: PMC4257651.

Bhuiyan MU et al 2014: Bhuiyan MU, Luby SP, Alamgir NI, Homaira N, Mamun AA, Khan JA, Abedin J, Sturm-Ramirez K, Gurley ES, Zaman RU, Alamgir AS, Rahman M, Widdowson MA, Azziz-Baumgartner E. Economic burden of influenza-associated hospitalizations and outpatient visits in Bangladesh during 2010. *Influenza Other Respir Viruses*. 2014 Jul;8(4):406-13. doi: 10.1111/irv.12254. Epub 2014 Apr 22. PMID: 24750586; PMCID: PMC4181799.

Appendix

Literature Review for effectiveness & safety

This literature search is an example of Level 1 review where intervention inputs taken from DCP3 or generated in an ad hoc manner (e.g., quick google search found one study of cervical cancer screening cost-effectiveness that was used to create an effectiveness parameter for that intervention).

Level of evidence of efficacy studies:

1. low (expert opinions, case series, reports, low-quality case control studies)
2. moderate (high quality case control studies, low quality cohort studies)
3. high (high quality cohort studies, individual RCTs)

4. very high (multiple RCTs, meta-analysis, systematic review, clinical practice guidelines).